



Chinese Pharmaceutical Association
Institute of Materia Medica, Chinese Academy of Medical Sciences

Acta Pharmaceutica Sinica B

www.elsevier.com/locate/apsb
www.sciencedirect.com



REVIEW

Intestinal barrier in chronic gut and liver diseases: Pathogenesis and therapeutic targets

Yongxin Zhang^{a,b}, Yameng Liu^{a,b,*}, Xinyu Liang^{a,c}, Yingquan Wen^{a,d},
Jingjie Zhao^{a,e,f}, Yong He^{a,b,*}, Qing Xie^{g,*}, Cen Xie^{a,b,e,g,*}

^aState Key Laboratory of Drug Research, Shanghai Institute of Materia Medica, Chinese Academy of Sciences, Shanghai 201203, China

^bUniversity of Chinese Academy of Sciences, Beijing 100049, China

^cDivision of Nephrology, Longhua Hospital, Shanghai University of Traditional Chinese Medicine, Shanghai 200032, China

^dSchool of Life Sciences and Technology, Tongji University, Shanghai, 200127, China

^eSchool of Life Science and Technology, ShanghaiTech University, Shanghai 201210, China

^fLingang Laboratory, Shanghai 200031, China

^gRuijin Hospital, Shanghai Jiao Tong University School of Medicine, Shanghai 200025, China

Received 19 January 2025; received in revised form 22 June 2025; accepted 29 August 2025

KEY WORDS

Intestinal barrier;
Barrier dysfunction;
Barrier integrity detection;
Factors affecting intestinal integrity;
Gut and liver diseases;
Therapeutic targeting barrier repair;
Tight junction;
Dysbiosis

Abstract The intestinal barrier is the primary defense that separates the host from the external environment, possessing several crucial physiological functions, including nutrient digestion, absorption, and protection against potentially harmful dietary antigens and pathogenic microorganisms. Nevertheless, various factors, such as diet, medications, circadian rhythm disturbances, gut microbiota, microbial metabolites, and genetic predisposition, can disrupt the intestinal barrier. Such disruption may lead to bacterial translocation, subsequently triggering enterohepatic and systemic inflammation. Impaired intestinal barrier has been implicated in the pathogenesis of numerous diseases, particularly chronic gut and liver diseases. In this review, we will summarize the fundamental functions of intestinal barrier and discuss clinical correlations between intestinal barrier dysfunction and diseases such as colitis, colorectal cancer, and chronic liver diseases including metabolic dysfunction-associated steatohepatitis, alcohol-associated liver disease, and primary sclerosing cholangitis. Additionally, we will also highlight some potential therapeutic strategies aimed at restoring barrier integrity to improve disease management.

*Corresponding authors.

E-mail addresses: yameng_liu@simm.ac.cn (Yameng Liu), heyong@simm.ac.cn (Yong He), xieqingrjh@163.com (Qing Xie), xiecen@simm.ac.cn (Cen Xie).

Peer review under the responsibility of Chinese Pharmaceutical Association and Institute of Materia Medica, Chinese Academy of Medical Sciences.

<https://doi.org/10.1016/j.apsb.2025.08.028>

2211-3835 © 2025 The Authors. Published by Elsevier B.V. on behalf of Chinese Pharmaceutical Association and Institute of Materia Medica, Chinese Academy of Medical Sciences. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

The preservation of barrier integrity is a crucial indicator of health and is essential for withstanding environmental stress while supporting physiological functions^{1–5}. This internal biological barrier facilitates the establishment of ventricular electrophysiological and chemical gradients, osmotic equilibrium, coordination of ventricular communication, and normal turnover of metabolic circuits. Meanwhile, the external barrier, primarily represented by the intestinal cavity, serves as a “boundary line” that prevents the translocation of harmful substances and microorganisms from the gut into the systemic circulation^{6,7}. The intestinal barrier is a complex structure and comprises several components: mucus layer, epithelial tight junctions (TJs) and adherens junctions (AJs), stem cell niches, mesenchymal cells, resident immune cell population, and symbiotic gut microbiota^{8–16}. The physical barrier is formed by a dense epithelial layer, composed of specialized cells originating from crypt stem cells, including enterocytes, goblet cells, Paneth cells, tuft cells, M cells, and other enteroendocrine cells^{17–21}. These cells are interconnected by TJs that regulate intestinal homeostasis, working in conjunction with biochemical barriers constructed by self-secreting host factors and microbial metabolites.

Over the past few decades, many clinical disorders have been associated with intestinal barrier dysfunction^{22–24}. These conditions include intestinal diseases, such as inflammatory bowel disease (IBD), which is comprised of ulcerative colitis (UC) and Crohn’s disease (CD), as well as irritable bowel syndrome (IBS) and colorectal cancer (CRC). Beyond the gut, chronic liver diseases such as metabolic dysfunction-associated steatotic liver disease (MASLD), alcoholic liver disease (ALD), and primary sclerosing cholangitis (PSC), and systemic metabolic disorders such as diabetes and obesity, are also linked to impaired intestinal barrier integrity. However, most clinical data remains correlational, leaving unresolved whether intestinal barrier damage is a cause or a consequence of these diseases²⁵. Currently, no US Food and Drug Administration (FDA)-approved medications are specifically designed to repair intestinal barrier damage. Most therapies focus on the prevention and management of the related diseases, often relying heavily on immunosuppressants to control inflammation^{26,27}. Unfortunately, persistent barrier impairment and delayed healing can reduce the efficacy of these treatments, and potentially lead to treatment tolerance and even relapse²⁸. Therefore, it is essential to develop therapeutic strategies that directly target the epithelial barrier. Further research is necessary to better understand the physiological regulation of the intestinal barrier, the mechanisms underlying its dysfunction, and its role in disease development and progression.

This review discusses the physiological structure and function of the intestinal epithelial barrier, the internal and external factors affecting its integrity, and the role of barrier impairment in disease pathogenesis. We aim to provide insights that may guide future research on developing therapies targeting the intestinal barrier dysfunction and related diseases.

2. Structural composition of intestinal barrier

The intestine is a unique organ with crucial roles in food digestion, nutrient absorption, dynamic host-environment interaction, and body homeostasis maintenance. To protect the host from external threats, such as prolonged exposure to dietary antigens and pathogenic microorganisms, intestinal epithelial cells (IECs) form multiple types of barriers, including a mechanical barrier, a mucus layer enriched with symbiotic microbes, and an immune barrier composed of immune cells and their active substances (Fig. 1). The mechanical barrier consists of a compact layer of IECs formed by TJs, which ensure the structural integrity of the intestine, regulate intestinal permeability, and control the transportation of water and macromolecules²⁹. The mucus layer, made of mucin secreted by goblet cells, provides both habitat and nutrients for symbiotic bacteria. Its unique structural characteristics restrict the penetration of pathogens, further enhancing the physical isolation function of TJs³⁰. Furthermore, there is a stem cell niche located at the base of the crypt, which consists of crypt-resident intestinal stem cells (ISCs), mesenchymal cells, immune cells, and intestinal secretory cells, such as goblet and Paneth cells. This niche is highly proliferative and thus is responsible for tissue renewal and intestinal barrier repair. It also mediates antigen phagocytosis and releases antimicrobial peptides (AMPs) to maintain internal homeostasis by eliminating potential pathogens³¹. Together, these elements create physical and biochemical barriers that protect the host and regulate communication between the internal and external environments. This intricate balance is essential for the proper functioning of the gastrointestinal system and the maintenance of overall bodily equilibrium.

2.1. Mucus layer composition and immune surveillance

The mucus layer serves as the first line of defense for the gastrointestinal tract, primarily consisting of 90%–95% water, 1%–2% lipids, and 1%–5% mucins. The mucins are highly glycosylated with 50%–80% carbohydrates (w/w) by glycosyltransferases. The diverse and complex structure of glycans in mucin provides colonization sites and nutrients to the mucus-associated bacteria that inhabit the mucus niche³². Mucin 2 (MUC2), the main mucin expressed in the gastrointestinal tract, is secreted by goblet cells upon bacterial exposure *via* meprin β -mediated cleavage, forming a protective mucin layer³³. The structure of the mucus layer allow symbiotic microbiota to colonize in the outer mucus layer and utilize their polysaccharide-degrading enzymes to derive nutritional energy from mucin *O*-glycan^{9,34,35}. This interaction helps regulate the structure and transcription of the proximal colon microbiota³⁶. A low-fiber diet has been shown to promote degradation of host mucins by the microbiota, leading to a thinner mucus layer and, consequently, a weakened barrier³⁷. *Muc2*-deficient mice displayed increased colon histological damage, greater bacterial translocation to the liver, and a marked reduction in intestinal TJ proteins³⁸. Furthermore, impairment of mucus barrier integrity and function due to mucin *O*-glycosylation disorders has been implicated in the




Figure 1 Structural composition of the intestinal barrier. The intestinal barrier is the largest barrier separating the host from the external environment, and has the physiological function of protecting the host from continuous antigen exposure to and invasion by pathogenic microorganisms. Maintaining the integrity of the intestinal barrier is important for overall health. The intestinal barrier comprises the mucus layer barrier, the mechanical barrier, the stem cell niche formed by intestinal stem cells and differentiated specialized cells, and the immune barrier. The mucus layer provides a habitat and nutrient components for the symbiotic microbiota, while limiting the penetration of pathogens due to its unique structural characteristics. The dense layer of epithelial cells formed by tight junctions determines the structural integrity of the intestine and intestinal permeability, which controls the transport of water and macromolecules. Crypt-resident stem cells maintain the continuous regeneration and damage repair of the intestinal epithelium, and mediate differentiation into different types of intestinal cells to support the functional diversity of the intestinal epithelium. The intestinal epithelial barrier is essential for preserving the integrity, function, and homeostasis of the intestinal structure. ZO-1, zonula occludens 1; MLCK, myosin light chain kinase; JAM, junctional adhesion molecule; MUC, mucin; AMP, antimicrobial peptides. Created with Biorender.com.

pathogenesis of metabolic diseases³⁹. Recent studies have identified Gasdermin D (GSDMD), a pore-forming effector protein involved in pyroptosis, regulates mucin secretion in goblet cells by promoting calcium-dependent exocytosis *via* scinderin-mediated F-actin depolymerization. GSDMD deficiency disrupts the mucus barrier, enabling pathogen adhesion to the epithelium and leading to the intestinal disease pathogenesis⁴⁰.

The viscosity of mucus also supports the retention and diffusion of antibacterial substances from Paneth cells and goblet cells, and immune-derived cytokines, which collectively protect the host from infection³². AMPs, such as crypt defensins, antimicrobials, and lysozymes, are produced by Paneth cells. These AMPs are abundantly on the surface of intestinal epithelium and capable of eliminating harmful microorganisms directly. In neonatal non-obese diabetic mouse models, colonic AMP deficiency due to ecological dysregulation contributes to pancreatic autoimmunity in type 1 diabetes⁴¹. AMPs, along with microbe-specific

immunoglobulin IgA, play a crucial role in maintaining barrier stability and inhibiting inflammation⁴². Their regulation is influenced by IL-17 and IL-22, which are produced by T helper 17 (Th17) cells and type III innate lymphoid cells (ILC3)⁴³. The latter relies on the dendritic cell (DC)-associated Mincle signaling pathway, which is linked to the tyrosine kinase-coupled C-type lectin receptor. In the absence of Mincle or when tyrosine kinase is impaired, the synthesis of intestinal regenerating islet-derived III- γ (RegIII γ) and IgA is reduced, permitting gut microbiota translocation that drives hepatic inflammation and lipid metabolism dysregulation⁴³. These findings demonstrate the crucial role of mucus layer integrity in maintaining gut–liver homeostasis.

2.2. Composition of epithelial junctions

Microvilli, apical projections of IECs, expand the intestinal surface area, optimizing nutrient contact with specific receptors and

digestive enzymes expression to enhance intestinal absorptive function⁴⁴. The densely packed microvilli form the brush border, a barrier that also prevents bacterial attachment to IECs. In patients with CD, studies revealed transcriptional dysregulation in genes associated with the brush border and reduced microvilli length, suggesting a possible link between microvilli abnormalities and epithelial lesions⁴⁵. The microvilli-stabilizing actin ACT-5, forming a bundle structure that maintains the normal morphology of intestinal microvilli, is disrupted by virulence factors of enterohemorrhagic *Escherichia coli* via CDK1-Formin signaling axis, leading to microvilli effacement, IEC barrier dysfunction, and worsens diseases such as severe diarrhea and hemorrhagic colitis⁴⁶.

The apical junction complex consists of TJs, AJs, and desmosomes. Under physiological conditions, TJs localize near the plasma membrane between adjacent cells, where anastomosis chains merge neighboring cell membranes and seal the paracellular space^{29,47}. One of the key roles of TJs is to form an epithelial permeability barrier that selectively regulates solute transport and charge flux, providing an essential “gate and fence” function⁴⁸. The claudin protein family, which has a four-transmembrane domain structure, is integral to TJ formation. Claudins determine the charge selectivity of the paracellular channel and are essential for proper TJ function⁴⁹. Generally, the downregulation of claudins promotes the progression of inflammatory and metabolic disorders due to barrier and paracellular channel defects. Conversely, elevated claudin-2 levels accelerate the onset of IBD, as claudin-2 acts as a pore barrier to increase intestinal cationic permeability, fostering inflammatory responses⁵⁰.

Zonula occludens 1 (ZO-1), the first discovered TJ protein, serves as a scaffold that positions several transmembrane proteins, such as occludin, claudin, and junctional adhesion molecule (JAM), at the cell–cell junctions to strengthen the bond between TJs and the actin cytoskeleton^{29,51}. ZO-1 anchors the TJ cytoskeleton, maintaining structural integrity and serving as a key barrier function marker. While essential for TJ stability, its direct role in TJ assembly and signaling remains undefined. Reduced ZO-1 expression in IBD might result from abnormal orientation of the mitotic spindle due to disrupted Wnt- β -catenin signaling, which inhibits epithelial proliferation and mucosal repair rather than impaired TJ-dependent barrier function⁵².

TJs are central to barrier regulation, and their breakdown can be a primary pathophysiological factor in pathogen invasion. Loss of TJs compromises the barrier and is associated with aberrant adenosine cAMP/protein kinase A (PKA)–cAMP response element-binding protein (CREB) signaling pathway⁵³. Additionally, the cytoskeleton critically regulates TJ dynamics and epithelial integrity. The pathogen *Chlamydia trachomatis* utilizes the effector protein *TepP* to disrupt host cytoskeleton-regulated tyrosine phosphorylation during the initial phase of infection, leading to TJ disassembly and further infection⁵⁴. TRIM40, an E3 ubiquitin ligase, disrupts epithelial barrier integrity by targeting ROCK1, a critical kinase for cell–cell junction integrity, destabilizing cortical actin and promoting IBD pathogenesis⁵⁵.

2.3. Stem cell niche and specialized epithelial cells

The continuous proliferative capacity of the intestinal crypt supports the maintenance and renewal of intestinal barrier. ISCs, located at the base of the crypt, plays a key role in both tissue repair and self-renewal to resist hazards within the digestive tract

by continually replenishing damaged IECs⁵⁶. In response to pathogen infection or epithelial injury, elevated IL-1R1 signaling induces the production of RSPO3, a Wnt agonist that promotes self-renewal of ISCs and facilitates IL-22-driven barrier repair⁵⁷. Intestinal barrier damage is often attributed to the dysfunction of stem cell niche, and several studies have demonstrated the environment factors, psychological stress, and aging are able to impair ISC lineage commitment. ISCs differentiate into diverse secretory and absorptive enterocytes in response to microenvironmental signals and nutrient availability⁵⁶. Psychological stress modifies sympathetic nerve output, which affects ISC mitochondrial bioenergetics via microbial metabolites and thus influences ISC differentiation and epithelial barrier repair. This mechanism partially explains the increased vulnerability to intestinal diseases in patients with mental disorders⁵⁸. Similarly, aging compromises barrier integrity via ISC depletion and dysbiosis, resulting from diminished Wnt signaling produced by Paneth and mesenchymal niche cells essential for ISC maintenance⁵⁹.

In addition to Paneth cells and goblet cells, tuft cells represent another specialized secretory IEC type within the epithelial barrier. Tuft cells play a protective role against pathogen infections by secreting the cytokine IL-25, which activates lamina propria lymphocytes to drive type 2 immunity and pathogen clearance⁶⁰. Tuft cells interact with pathogen metabolites via membrane receptors like Vmn2r26, initiating G-protein-coupled signaling to produce prostaglandin D2 (PGD2). This stimulates goblet cell mucus secretion and SpiB-mediated tuft cell proliferation, amplifying antimicrobial defense⁶¹. Furthermore, tuft cells can function as reserve ISCs during intestinal injury to aid the barrier repair⁶².

3. Factors in the regulation of intestinal epithelial barrier

The integrity of the intestinal barrier is affected by a variety of intrinsic and extrinsic factors. Barrier dysfunction can be triggered by genetic susceptibility, the preference for a western diet, the use of antibiotics and drugs, disruptions in circadian rhythms, psychological stress, and aging (Table 1)^{37,64–70,72,73,89,95–97,99,106,108,113,118,133,134,136–138,140,146–148,151–153,209}. These influences often lead to imbalances in gut microbiota, which can induce systemic metabolic disorders (Fig. 2). The subsequent inflammation due to these metabolic disturbances further weakens intestinal barrier, and thus establishes a damaging cycle of ongoing deterioration.

3.1. Genetic susceptibility

Advancements in genome-wide association studies (GWAS) have notably enhanced the identification of disease-susceptibility genes and the understanding of associated biological pathways, which are valuable for clinical translation. These studies analyze genetic variants in genome to explore the relationships between genotype and phenotype⁶³. While direct links between genetic mutations and intestinal barrier dysfunction are limited, GWAS has identified numerous loci associated with IBD development^{64,65}. For instance, the *IL10* gene mutation was one of the first mutations found to induce IBD⁶⁶. *IL10* knockout mice spontaneously develop colitis and increase intestinal microbiota translocation. Mutations in the IL10 receptor (IL10R), composed of IL10R1 and IL10R2 proteins encoded by *IL10RA* and *IL10RB* genes, have been associated with early-onset enterocolitis. These mutations impair IL10-induced signaling, that potentially increases the secretion of

Table 1 Factors and mechanisms of gut and liver diseases linked to intestinal barrier dysfunction.

Class	Type of disease	Affecting factors	Potential mechanism
Gut disease	UC and CD	Genetic predisposition ^{64–68,70,137}	Defects in genes and pathways associated with innate and adaptive immunity, autophagy, and mesenchymal cells may mediate CD susceptibility ⁶⁴ . Mutations in IL10R induce hyperinflammatory immune responses in the intestine ⁶⁶ . Defect in <i>SLC26A3</i> , a novel susceptibility gene for IBD, causes abnormal expression of TJ and AJ through a post-transcriptional mechanism ⁶⁹ . <i>ST6GALNAC1</i> (<i>ST6</i>) mutation leads to abnormal mucin glycosylation modification, which causes dysbiosis and inflammatory activation by impairing the integrity and function of the intestinal mucus barrier ⁷⁰ . Impaired cellular adhesion regulation during intestinal wound repair and disease development ¹³⁶ .
		BA dysregulation ^{72,73}	Cholic acid impede <i>Lgr5</i> ⁺ ISC proliferation by suppressing PPAR α , ultimately impairing epithelial barrier regeneration ⁷² . TGR5-mediated BA signaling in ISCs activates SRC/YAP-dependent transcriptional programs driving epithelial renewal ⁷³ .
		Dysbiosis ⁹⁶	Depletion of SCFA-producing microbiota leads to barrier dysfunction <i>via</i> suppressing TJ related protein expression and mucus production ^{96,97} .
		Circadian rhythms ⁹⁹	Disruption of this diurnally regulated diet increases the burden of exposure to dietary antigens and microbial stimuli through the disruption of microbiome-MHC II-IL-10 epithelial barrier axis, driving Crohn-like enteritis ⁹⁹ .
		Drug usage ^{89,118}	Antibiotics disrupted mucin secretion, enabling microbiota penetration and translocation into circulation, thereby causing intestinal inflammation ¹¹³ . Anti- PD-1 immunotherapy reduces IL-22 production by LT α cells, thereby exacerbating intestinal inflammation ¹¹⁸ .
		Diet ^{37,106}	Transient HFD exacerbates colitis through elevating intestinal BAs and TNF levels, which together trigger cytotoxic BA accumulation in the endoplasmic reticulum, ultimately inducing intestinal epithelial apoptosis <i>via</i> the IRE1 α /XBP1 pathway ¹⁰⁶ . Fiber-deficient diets shift microbial metabolism toward mucin degradation, compromising the mucus barrier and creating an ecological niche for pathobiont expansion ³⁷ .
		Mucosal immune activation	TNF- α signaling activates the MLCK promoter, increasing long-chain MLCK expression ¹³³ . IL-13 affects epithelial barrier permeability by increasing the expression of cladin-2, and causes epithelial cell apoptosis and barrier resitutation velocity ¹³⁴ .
	IBS CRC	Genetic predisposition ¹³⁸	
		Dysbiosis ⁹⁵	FFAR2-mediated microbial metabolite sensing orchestrates protective effects on epithelial integrity, microbial ecology, and immune surveillance within the tumor microenvironment ⁹⁵ .
		Diet ¹⁰⁸	HFD-induced colorectal carcinogenesis involves synergistic effects of gut dysbiosis, microbial metabolite LPA-mediated CRC cell proliferation and impair cell junction, and compromised epithelial barrier function ¹⁰⁸ .
Chronic liver diseases	MASLD	Genetic predisposition ¹⁴⁰	Early genetic events disrupt intestinal barrier-related protein and results in microbial product infiltration into adenomatous tissue, triggering IL-23/IL-17-mediated tumour growth ¹⁴⁰ .
		Dysbiosis ¹⁴⁶	MASH patient fecal EVs drive gut–liver axis dysfunction through concurrent TJ suppression, TLR4/LPS-mediated inflammation, and HSCs activation ¹⁴⁷ .
	ALD	Alcohol exposure	ALDH2 deficiency exacerbates alcohol-induced oxidative stress, intestinal epithelial apoptosis, and TJ/AJ degradation, thereby increasing gut permeability and endotoxemia to drive systemic inflammation and liver injury ¹⁴⁸ .
		Dysbiosis ¹⁵¹	Alcohol depletes intestinal cDCs, impairing IL-12/IFN- γ -mediated AMP production and reducing <i>A. muciniphila</i> abundance ²⁰⁹ . <i>Klebsiella pneumoniae</i> destroys the epithelial barrier, initiates bacterial translocation and the Th17 immune response in the liver, and increases the susceptibility to hepatobiliary injury ¹⁵¹ .
	PSC	Dysbiosis ¹⁵¹	LPS-mediated hepatocellular NF- κ B activation protects cholestatic liver injury through negative feedback by inhibiting BA metabolism ¹⁵³ .
		ISCs stemness impairment	BDL-induced cholestatic injury involves the activation of ER stress and disruption of ISCs stemness that exacerbate systemic inflammation ¹⁵² .




Figure 2 Factors affecting intestinal permeability and barrier function. The integrity of the intestinal barrier can be compromised by several factors, including genetic predisposition, western dietary preferences, use of antibiotics and other drugs, circadian rhythm disruption, and other internal and external factors. Inherited traits that may influence the structure and function of the intestinal barrier. Western dietary habits, characterized by high fat and sugar intake, can negatively impact gut health. Use of antibiotics and other medications can disrupt the balance of gut microbiota and compromise barrier function. Circadian rhythm disruption induced by irregular sleep patterns and unhealthy lifestyle factors can also influence gut health and permeability. Moreover, stress, infections, and environmental toxins can also play a role in degrading barrier integrity. The impairment of intestinal barrier can lead to microbial imbalance and a pro-inflammatory microenvironment. The resulting inflammation can further exacerbate intestinal barrier damage, creating a vicious cycle that contributes to the progression of various metabolic diseases. Created with Biorender.com.

TNF- α and other pro-inflammatory cytokines, thus exacerbating inflammation and weakening intestinal tolerance⁶⁶. GWAS has also shown the significant downregulation of the *SLC26A3* gene, which encodes the DRA protein, an intestinal chloride anion transporter. This gene ranks among the top 1% of harmful IBD variants in the human genome^{67,68}. Reduced expression of *SLC26A3* significantly increases colonic paracellular permeability, lowers the expression of TJ and AJ proteins, and thus heightens susceptibility to IBD⁶⁹.

Additionally, glycoproteomic profiling has revealed that patients with congenital IBD may carry mutations in the *ST6GALNAC1* (*ST6*) gene, which encodes a sialyltransferase crucial for maintaining mucus barrier homeostasis. Mutations in this gene lead to a reduction in the thickness of the intestinal mucosal layer and disruption of its protective function⁷⁰. Elucidating these regulatory mechanisms is important for comprehending the pathogenesis of congenital IBD. Furthermore, barrier function is also influenced by

interactions of genetic predisposition and environmental factors. Mutations in DNA methyltransferase coding gene *DNMT3A*, which is associated with an increased risk of IBD, have been shown to reduce goblet cell number, shorten AJ complex, and increase intestinal permeability⁷¹. These changes increase susceptibility to colitis and hinder epithelial regeneration and repair processes.

3.2. Bile acids

Bile acids (BAs), which are the end products of cholesterol metabolism, act as natural emulsifiers that facilitate the absorption of intestinal lipids and the biliary excretion of cholesterol and phospholipids. Beyond these traditional functions, BAs also serve as hormones that regulate systemic metabolism and immune responses by binding to receptors such as Farnesoid X Receptor (FXR) and Takeda G-protein-coupled receptor 5 (TGR5). Recent studies emphasize the critical role of BAs in maintaining the

intestinal epithelial barrier. BA signals stimulate self-renewal of *Lgr5*⁺ ISCs for intestinal barrier repair and homeostasis maintenance. However, excessive accumulation of the primary BA cholic acid in the intestine can impede *Lgr5*⁺ ISC proliferation by hindering fatty acid oxidation *via* peroxisome proliferator-activated receptor (PPAR α), thereby decelerating epithelial barrier repair⁷². Similarly, TGR5-mediated-signaling plays a vital role in preserving metabolic homeostasis, especially in the intestine where its expression is notably high. Activated by its natural ligand deoxycholic acid (DCA), a secondary BA metabolized from cholic acid, the TGR5 in ISCs triggers the transcriptional activation of downstream factors such as Yes-associated protein 1 (YAP1) and its regulator SRC, which collectively drive intestinal epithelial regeneration⁷³.

Furthermore, pregnane X receptor (PXR), a key bile acid-sensing nuclear receptor, modulates intestinal epithelial barrier homeostasis through its dual regulation of xenobiotic metabolism and innate immune regulation. Clinical studies have demonstrated significant downregulation of PXR and its downstream target genes in UC patients, suggesting that impaired PXR-mediated responses may contribute to defective barrier repair following intestinal injury⁷⁴. PXR-deficient mice exhibit severe intestinal epithelial dysfunction and heightened susceptibility to multiple intestinal injury models⁷⁵. The endogenous PXR ligand lithocholic acid (LCA) significantly attenuates pro-inflammatory signaling in necrotizing enterocolitis⁷⁶. PXR activation mediates these protective effects by antagonizing the NF- κ B pathway, suppressing pro-inflammatory cytokine release, and preserving TJ integrity during inflammation^{77–79}. Bile acids function as endogenous ligands for the vitamin D receptor (VDR), exerting protective effects particularly in intestinal homeostasis⁸⁰. VDR activation plays a crucial role in maintaining intestinal barrier integrity by regulating epithelial differentiation and enhancing TJ expression^{81–83}. Furthermore, VDR signaling is essential for supporting Paneth cell antimicrobial function⁸⁴. However, this protective mechanism becomes impaired during the pathogenesis of IBD and CRC. Clinically, dysregulation of the VDR signaling pathway is strongly associated with accelerated CRC progression and unfavorable clinical outcomes. Beyond its direct anti-proliferative effects on tumor cells, VDR activation contributes to CRC suppression by enhancing mucosal barrier integrity, thereby restricting tumor advancement^{85,86}.

3.3. Gut microbiota and derived metabolites

Under normal physiological conditions, the gut microbiota exists symbiotically with the host, primarily confined to the outer mucus layer by the intestinal epithelial barrier. Nevertheless, some microorganisms have evolved mechanisms to bypass this barrier, compromising the structural and functional integrity of the gastrointestinal tract. Although antibiotics might initially seem protective, their use is often counterproductive, as they can weaken the barrier function and increase the risk of new-onset IBD^{87,88}. A population-based study in Sweden revealed a correlation between cumulative systemic antibiotic exposure and a higher risk of new-onset IBD⁸⁹. In contrast, the gut microbiota has positive effects on intestinal barrier repair. By comparing with germ-free mice, the presence of gut microbiota promotes the growth of colon epithelial precursor cells *via* the signaling of activated macrophages within the stem cell niche to aid the restoration of damaged epithelial barriers in conventional mice, thus confirming their role in barrier repair⁹⁰.

The gut microbiota exerts their protective effects on the integrity of intestinal barrier by producing metabolites, including secondary BAs, which influence the function the intestinal barrier to impact overall gut health. In addition to BAs, clinical evidence has shown that the levels of another microbial metabolite tryptophan are significantly decreased in serum of patients with IBD⁹¹. Further studies suggest that alterations in the tryptophan metabolism are pivotal to disease progression and prognosis^{92,93}. Tryptophan metabolites, specifically indole derivatives, serve as natural ligands for the aryl hydrocarbon receptor (AhR), which is widely expressed in the gut. AhR activation regulates regulatory T cells (Tregs) and Th17 cells, as well as the production of their cytokines, which are important for barrier protection, repair, and homeostasis maintenance, with IL-22 being a key effector. The kynurenine (Kyn) pathway, a major tryptophan metabolism route, is correlated with IBD progression⁹¹. Indoleamine 2,3-dioxygenase (IDO), a rate-limiting enzyme in the Kyn pathway, promotes tryptophan metabolism to Kyn pathway under inflammatory conditions. IDO deficiency or inhibition can alleviate intestinal endotoxemia and chronic inflammation, as well as enhance intestinal mucosal and immune barriers, particularly in obesity models⁹⁴. In addition to AHR activation, these metabolites enhance barrier integrity *via* PXR-mediated mechanisms. Specifically, indole-3-propionic acid (IPA) serves as an endogenous PXR agonist, attenuating intestinal permeability and inflammatory responses through TLR4 signaling⁷⁵. Previous studies reveal that PXR activation preserves intestinal barrier integrity through coordinated mechanisms involving stabilization of TJ protein ZO-1 localization, suppression of myosin light chain kinase (MLCK) expression, and inhibition of JNK1/2 phosphorylation⁷⁹. In summary, tryptophan-derived indole metabolites mediate dual protective effects on the intestinal epithelium through simultaneous activation of both PXR and AHR signaling pathways.

Short-chain fatty acids (SCFAs), another essential group of microbiota-derived metabolites, also support intestinal barrier function. SCFAs, such as butyrate, acetate, and propionate, are produced by the breakdown of dietary fiber by symbiotic flora, such as Firmicutes. These SCFAs act on the GPCRs, such as GPR43 or free fatty acid receptor 2 (FFAR2), to enhance the integrity of the intestinal barrier and protect CD8⁺ T cells and DC cells from exhaustion and overactivation, thereby preventing the CRC development⁹⁵. Butyrate, in particular, is known to inhibit histone deacetylases (HDACs) in the intestine to regulate gene transcription and aid in barrier restoration by modulating the actin-binding protein synaptopodin (SYNPO) at the TJ of colon epithelium and F-actin stress fibers⁹⁶. GPCR signaling and HDAC inhibition are suggested to work together to maintain the integrity of the epithelial barrier. Furthermore, SCFAs stimulate goblet cells to secrete mucins, which in turn promote the growth of mucin-degrading bacteria to facilitate ISC differentiation into secretory lineages by consuming mucins, thereby ensuring a robust regenerative capacity for the epithelium⁹⁷.

3.4. Circadian rhythms

Circadian rhythms, often referred as the biological clock, are an internal control system for metabolic homeostasis, evolved to synchronize periodic environmental signals such as the light–dark cycle. This system orchestrates gene expression at specific times through transcriptional, post-transcriptional, and post-translational modifications to coordinate the optimal metabolic adaptation⁹⁸. Several studies have demonstrated a strong correlation between

circadian disruptions and metabolic disorders. Specifically, irregular feeding schedules increase the exposure of small intestine to dietary antigens and microbial stimuli, resulting in a greater burden on the major histocompatibility complex II (MHC II) of IECs. Such disruption weakens the regulatory function of the intestinal microecology and diminish IL-10 secretion⁹⁹. Circadian rhythms also impact neuroimmune circuits in the gut, which are activated by feeding behavior and can lead to pathological changes under dietary irregularities. Food intake triggers intestinal neurons to secrete vasoactive intestinal peptide (VIP), which upregulates proteins related to lipid absorption and transport. Concurrently, VIP reduces AMP levels in IECs and decreases the production of IL22 by ILC3¹⁰⁰. Such disruptions in dietary rhythms facilitate intestinal colonization of pathogens, particularly when the barrier function of neuroimmune circuits is compromised¹⁰⁰.

Gut microbiota itself also exhibits circadian fluctuations in abundance and function, referred to as microbial oscillators, which impact the host circadian rhythm through microbial metabolites or autoantigens^{101,102}. This interplay between the gut microbiota and host biological clock significantly influences barrier integrity and innate immune responses. For instance, SCFAs help regulate the circadian phase shift in the small intestine by inhibiting HDAC activity¹⁰³. Segmented filamentous bacteria (SFB) in the gut drive rhythmic ILC3 circuit oscillations, in sync with host rhythm, thereby mediating diurnal alterations in infection resistance *via* time-dependent expression of AMPs¹⁰⁴.

3.5. Diet

The western diet, high in fat and sugar, is closely associated with the increased prevalence of metabolic diseases in western societies¹⁰⁵. Dietary components significantly shape the gut microenvironment by altering the gut microbiome and the production of secondary metabolites that actively participate in host physiological processes¹⁰⁶. Studies have shown that shifts in gut microbiota induced by the western diet raise DCA levels, which impair Paneth cells through activation of the intestinal FXR and type I interferon (IFN) signaling pathways¹⁰⁷. Animal studies have further revealed that high-fat diet (HFD) exacerbates intestinal barrier damage across different colon cancer models, including the azomethane (AOM)-dextran sulfate sodium (DSS)-induced model as well as the *Apc*-mutation-induced spontaneous model. The HFD-compromised gut barrier allows more pathogens and derived metabolites to penetrate the epithelium, thereby accelerating CRC development¹⁰⁸.

In addition to high-fat intake, excessive dietary fructose consumption can also harm the epithelial barrier. Elevated fructose levels increase circulating endotoxins and thus activate toll-like receptor 4 (TLR4) on macrophages to trigger systemic inflammatory responses¹⁰⁹. Glucose metabolism, meanwhile, presents both therapeutic potential for metabolic syndrome and a key coordinator of intestinal barrier function. In mouse models of type 2 diabetes mellitus (T2DM) with leptin deficiency (*ob/ob*) and leptin receptor (*LepR*) deficiency (*db/db*), hyperglycemia disrupts the intestinal barrier by inducing glucose transporter 2 (GLUT2)-dependent transcriptional reprogramming in IECs, which subsequently impairs the integrity of TJ and AJ structures¹¹⁰.

3.6. Drug

The integrity of the gut barrier is intricately tied to the gut microbiome, with antibiotic usage, whether direct or indirect,

profoundly affecting the abundance and structure of microbiota. The consequences of antibiotic therapy on host health arise from potential loss of immunomodulatory microbiota ligands, alteration in secondary metabolites, and signals related to specific microbiota¹¹¹. Rifaximin- α , an antibiotic prescribed to prevent hepatic encephalopathy, safeguards the protective mucus membrane by reducing the population of sialidase-rich bacteria that break down mucus and increase levels of TNF- α and IL-17 in the gut microenvironment, which increases resistance to pathogens^{112,113}. In addition, rifaximin- α treatment ameliorates stress-induced intestinal barrier dysfunction by enriching *Lactobacillus* levels in the ileum¹¹⁴. However, antibiotic treatment can also reduce gut microbiota diversity, causing immune dysregulation and increased susceptibility to infections. In healthy adult mice treated with a broad-spectrum antibiotic cocktail, dysbiosis occurred along with impaired epithelial TJ integrity, as indicated by reduced ZO-1 expression and activation of the NLRP3 inflammasome⁸⁷.

Non-antibiotic drugs can also disrupt the intestinal barrier. Aspirin, widely used for its anti-inflammatory and analgesic properties, is associated with gastrointestinal damage. It activates intestinal FXR signaling and decreases the abundance of *Parabacteroides goldsteinii*, a bacterium that produces 7-keto-LCA, which inhibits the stemness of ISC and slows barrier repair¹¹⁵. Chemotherapies are another major contributor to gut barrier damage. For instance, 5-fluorouracil accelerates mucosal cell death over regeneration. Consequently, more than 40% of chemotherapy patients experience gastrointestinal injuries, manifesting as symptoms such as diarrhea, constipation, and indigestion¹¹⁶. Immune checkpoint inhibitors (ICIs) that target programmed death receptor 1 (PD-1) have revolutionized anti-tumor therapy. Although these antibodies re-establish regular immune response by blocking the PD-1 pathway, they often cause gastrointestinal toxicities¹¹⁷. PD-1 signaling has been found essential for regulating colon lymphoid tissue inducer (LTi) cells, a specific subgroup of ILC3, which are crucial for maintaining immune homeostasis. Loss of PD-1 signaling results in the over-activation of fatty acid oxidation in LTi cells, and feedback inhibits both LTi cell activation and IL-22 production. This imbalance leads to dysbiosis, intestinal barrier damage, and increased susceptibility to enteritis¹¹⁸.

4. Detection and analytical approaches of intestinal barrier integrity

In vivo assessment of intestinal barrier integrity can be performed through direct measurement of intestinal integrity, indirect analysis of biomarkers, and novel multi-omics-based analytical platforms. The assessment of intestinal permeability through urinary excretion of orally administered probe molecules represents a direct and widely utilized method. Typically, non-metabolized, poorly absorbed sugar probes, such as mannitol, rhamnose, sucralose, and lactulose, or low molecular weight polyethylene glycol (PEG) and ethylenediamine tetra-acetic acid (EDTA), are employed to evaluate intestinal barrier function¹¹⁹. To minimize potential interference from dietary sources, these probes are often isotope-labeled. Following oral administration, their urinary excretion rates are quantitatively assessed, serving as a reliable indicator of intestinal paracellular permeability¹²⁰. The lactulose-mannitol test is a widely employed clinical assay for assessing intestinal barrier integrity. In this test, the urinary excretion of the large molecular weight lactulose serves as an indicator of leaky

gut, reflecting the extent of intestinal barrier dysfunction at sites of epithelial damage. In contrast, the small molecular weight mannitol is predominantly absorbed *via* transcellular pathways, representing non-specific epithelial transport¹²¹. Consequently, the lactulose-to-mannitol ratio (LMR) is a well-established biomarker for evaluating intestinal permeability^{122,123}. In animal models, the absorption of FITC-labeled dextrans or chromium-51-labeled EDTA (⁵¹Cr-EDTA) into systemic circulation is commonly used to assess intestinal barrier integrity^{124,125} (Fig. 3)^{1,55,119,120,122,125,126,128–131}.

Furthermore, histopathological evaluation of endoscopic biopsy specimens allows for detailed examination of epithelial villous morphology and assessment of intestinal barrier integrity. Structural alterations in mucosal components can be systematically analyzed using mucin-specific Alcian Blue/Periodic Acid-Schiff (AB/PAS) staining for mucin layer evaluation, electron microscopy

for ultrastructural analysis, and immunofluorescence techniques for precise localization and quantification of TJ protein expression.

Indirect assessment of intestinal barrier integrity utilizes a comprehensive biomarker approach combining serum diamine oxidase (DAO) levels as an indicator of mucosal injury, D-lactate measurement for permeability evaluation, bacterial endotoxin such as LPS detection to assess microbial translocation, and fecal calprotectin (FCP), lipocalin-2 (LCN2), and albumin quantification for monitoring inflammatory activity^{122,126–131}. This approach is further enhanced by microbiome analysis, which provides systematic evaluation of the interconnected pathological features including mucosal damage, barrier dysfunction, and gut dysbiosis¹²². Recent advances in serum proteomics and genetic screening techniques have expanded the potential for early detection of barrier dysfunction while simultaneously supporting mechanistic investigations into its underlying causes^{1,55}.




Figure 3 Measurements of intestinal barrier integrity. *In vivo* assessment of intestinal barrier integrity can be performed through direct measurement of intestinal integrity, indirect analysis of biomarkers, and novel multi-omics-based analytical platforms. Direct assessment of intestinal permeability employs established methodologies including urinary excretion quantification of orally administered probe molecules such as lactulose and mannitol, intestinal leakage evaluation *via* the lactulose-to-mannitol ratio (LMR), and systemic absorption measurement using chromium-51-labeled ethylene diamine tetraacetate (⁵¹Cr-EDTA)^{119,120,122,125}. Furthermore, histopathological examination of endoscopic biopsy specimens enables direct evaluation of mucosal architecture, including villi morphology and the spatial distribution and expression patterns of TJ proteins. Indirect assessment of intestinal barrier integrity utilizes a comprehensive biomarker approach combining serum diamine oxidase (DAO) levels as an indicator of mucosal injury¹²⁶, D-lactate measurement for permeability evaluation¹²⁸, bacterial endotoxin such as LPS detection to assess microbial translocation¹³¹, and fecal calprotectin (FCP)¹²², LCN2¹²⁹, and albumin¹³⁰ quantification for monitoring inflammatory activity. This approach is further enhanced by microbiome analysis¹²², which provides systematic evaluation of the interconnected pathological features including mucosal damage, barrier dysfunction, and microbial dysbiosis. Recent advances in serum proteomics¹ and genetic screening techniques⁵⁵ have expanded the potential for early detection of barrier dysfunction while simultaneously supporting mechanistic investigations into its underlying causes. Created with Biorender.com.

5. Abnormal intestinal barrier function in the pathogenesis of gut and liver diseases

Dysregulation of the intestinal epithelial barrier is significant relevance to the development and progression of various gut and liver diseases. TJ dysfunction in the intestinal epithelium increases permeability, which facilitates the translocation of bacteria and their pathogenic factors across the barrier into systemic circulation and other tissues. This incursion triggers systemic inflammation and disrupts metabolic homeostasis, which are pivotal events in disease initiation and progression within the gut and liver. Although it remains uncertain whether the impairment of the intestinal barrier is a primary driver or a coincidental pathological feature, accumulating evidence suggests that the integrity of the intestinal barrier plays a critical role in the etiology of these diseases. A deeper understanding of the intrinsic and extrinsic factors that influence barrier function is necessary for unraveling the complex relationship between the intestinal barrier and disease pathology. This knowledge will provide directions for future research, leading to the identification of novel therapeutic targets that improve barrier function and address the root causes of gut and liver diseases.

5.1. Inflammatory bowel disease and colorectal cancer

IBD, including CD and UC, is characterized by chronic gastrointestinal inflammation, where epithelial barrier dysfunction is a key factor in disease onset and progression. Clinical studies show that altered intestinal TJ structures in UC and CD patients contribute to increased intestinal permeability¹³². Moreover, mucosal immune activation also leads to the production of cytokines such as TNF- α , IL-13 and IL-17. TNF- α induces epithelial barrier dysregulation by activating myosin light chain kinase (MLCK), which is considered a common regulatory pathway of acute TJ response to various immune and infectious stimuli¹³³. IL-13 reduces barrier function and increases claudin-2 expression, which allows selective entry of solutes and ions, along with epithelial cell apoptosis, ultimately leading to TJ breakdown^{50,134,135}. Further investigation suggests that increased paracellular permeability due to TJ dysfunction and cytoskeletal changes may precede the onset of disease, as seen in relatives of CD patients, who demonstrate early intestinal barrier irregularities and a higher prevalence of IBS^{136–138}. Similarly, large-scale cohorts have identified markers of barrier impairment up to three years before CD diagnosis. This supports that barrier dysfunction is a primary cause in disease progression rather than a secondary effect¹²³.

Moreover, epithelial barrier defects expose the intestinal lining to mutagenic compounds or prolonged inflammatory stimuli, which may initiate and promote colorectal tumorigenesis through mechanisms such as oxidative stress¹³⁹. In both early and advanced CRC, the expression of various barrier-related proteins is significantly reduced, which facilitates microbial translocation, triggers inflammation, and further accelerates tumor growth¹⁴⁰. However, the precise mechanisms underlying the abnormal expression of epithelial barrier components during the initiation and development of CRC remain unclear. Indeed, experimental models show that mice with compromised epithelial barriers are more susceptible to colitis and inflammation-driven tumors. For instance, the loss of Golgi membrane protein 1 (GOLM1) over-activates the Notch pathway, and thus disrupts intestinal homeostasis, alters in the differentiation and secretory cell profile of IECs, and decreases the number of goblet cells, all of which

increase intestinal permeability and pro-tumorigenic inflammation¹⁴¹. Mechanical barriers are also crucial in preventing CRC development. The tumor suppressor gene *N-myc* downstream-regulated gene 2 (*NDRG2*) has been shown to enhance the interaction between the E3 ligase FBXO11 and the E-cadherin inhibitor Snail, promoting Snail ubiquitination. This stabilization of E-cadherin strengthens AJs, thus limiting the development of colitis-associated tumors¹⁴².

5.2. Chronic liver diseases

Chronic liver diseases, including MASLD, ALD, and PSC, are frequently encountered with compromised intestinal barriers and dysregulation of the intestine–liver axis¹⁴³. Gastrointestinal substances, such as nutrients, secreted factors, and microbial metabolites, enter the liver *via* the portal vein and are metabolized for systemic utilization. This process renders the liver highly vulnerable to pathogen-associated molecular patterns (PAMPs), and makes it a primary target for enterogenic substances. In MASLD, endotoxemia is commonly observed, supporting the link between gut barrier integrity and liver inflammation. Similarly, DSS-induced gut barrier damage in mice leads to increased expression of pro-inflammatory genes and histological liver damage¹⁴⁴. In western diet-fed, DSS-treated *Il-10* knockout mice, T-cell infiltration in terminal ileums coincides with decreased BA levels in the portal vein and liver. This BA reduction suppressed liver FXR signaling to exacerbate MASLD pathogenesis¹⁴⁵. Gut dysbiosis, which compromises both the mucus layer and TJ integrity, has been identified as an early driver of MASH, as evidenced by metagenomic studies showing that gut microbiota dysregulation precedes liver disease onset in MASLD patients¹⁴⁶. Moreover, previous studies revealed that fecal extracellular vesicles from MASH patients could decrease TJ expression *via* a non-myosin light chain kinase (nmMLCK)-mediated mechanism, aggravating intestinal barrier dysfunction¹⁴⁷. Restoring the intestinal barrier in these cases can reverse liver lipid metabolism imbalances driven by colitis, particularly through the modulation of the secondary BA-TGR5/mTOR/oxidative phosphorylation pathway¹⁴⁵.

Additionally, alcohol exposure directly disrupts the integrity of the epithelial junction, and induces breakdown and dysfunction of intestinal barrier. Research demonstrates that the deficiency and inhibition of mitochondrial aldehyde dehydrogenase 2 (ALDH2), a key enzyme in alcohol metabolism, accelerate the binge drinking-induced degradation of intestinal TJ and AJ proteins, worsening intestinal epithelial dysfunction. Therefore, intestinal ALDH2 might be a potential target for alcohol-induced damage to the enterohepatic axis¹⁴⁸. Chronic alcohol intake also decreases conventional type 1 dendritic cells (cDCs) in the intestine, but not in the liver. This consequently leads to the downregulation of AMP production, a decline in protective *A. muciniphila* abundance, the breakdown of AJs, and ultimately causes liver injury *via* the IL-12-IFN γ signaling pathway¹⁴⁹.

PSC is a chronic cholestatic and inflammatory liver disease, with 70%–80% of cases co-occurring with UC. PSC patients exhibit distinct gut microbiota profiles compared to healthy individuals. Moreover, the presence of microorganisms in the portal vein suggests intestinal barrier disruption, which may allow for bacterial translocation and the influx of microbial products into the liver¹⁵⁰. Specific pathogenic bacteria, such as *Klebsiella pneumoniae* which are frequently detected in PSC patients, can induce the formation of monolayer epithelial pores through

contact-dependent apoptosis. This process lead to intestinal barrier damage, bacterial translocation, and activation of hepatic Th17 cells, which exacerbates inflammation and increases the vulnerability to liver and biliary injury¹⁵¹. Furthermore, in a murine model of cholestatic liver disease induced by bile duct ligation (BDL), intestinal barrier disruption occurs independently of dysbiosis. Knockout of C/EBP homologous protein (CHOP), an apoptosis-promoting transcription factor regulated by ER stress, significantly alleviates intestinal barrier damage, loss of ISC stemness, as well as liver inflammation and fibrosis in BDL mice. This finding indicates the important role of ER stress-mediated gut–liver axis imbalance and barrier dysfunction in cholestatic liver injury¹⁵². Interestingly, in a chronic IBD model, damage to the intestinal barrier stimulates a protective negative feedback circuit in PSC, by which the endotoxin lipopolysaccharides activate the NF- κ B pathway to suppress BA metabolism in hepatocytes, and then mitigates the progression of cholestatic liver disease¹⁵³. Therapeutic interventions, such as antibiotics¹⁵¹ or pan-caspase inhibitor IDN-7314¹⁵⁴, which attenuate dysbiosis-induced endotoxin translocation or NLRP3 inflammasome activation, have shown promise in treating PSC.

6. Potential interventions targeting the intestinal barrier

Immune-targeted therapies, including anti-inflammatory agents, immunosuppressants, and biologics, have proven effective in controlling inflammation in various intestinal diseases such as IBD. However, these treatments do not directly address the underlying damage to the intestinal barrier. Unfortunately, prolonged use of these medications may even increase the risk of infections, drug resistance, and disease recurrence, and serious adverse reactions, such as malignant tumors and death. Indeed, while reducing inflammation is crucial, promoting the healing and restoration of the epithelial barrier is pivotal for achieving long-term remission in intestinal disorders, especially IBD. Currently, no clinically approved therapies specifically target the repair and maintenance of the intestinal barrier. Given the close link between barrier dysfunction and the pathogenesis of both intestinal and liver diseases, timely repair and restoration of the epithelial barrier present a promising therapeutic strategy. Instead of focusing solely on symptom management, treatments that enhance epithelial healing and integrity could mitigate the progression of gut and liver diseases. Here, we summarized key therapeutic approaches for intestinal barrier restoration by targeting TJ adjustment, ecological regulation, ISCs regeneration, and immune regulation with supporting evidence from relevant clinical trials related to intestinal barrier integrity (Table 2). By strengthening the barrier, such therapies may also reduce bacterial translocation, inflammation, and the exacerbation of liver damage (Fig. 4)^{72,73,97,157,158,162,169,170,174,180,183–186,188}.

6.1. Tight junction adjustment

The maintenance of the intestinal barrier significantly relies on various TJ proteins. MLCK is considered a promising candidate target for developing barrier therapeutics for IBD due to its central role in regulating TJs and their breakdown. Deletion or inhibition of MLCK can effectively protect the barrier function from immune-induced damage, but these interventions do not prevent advanced colitis, which involves apoptosis and mucosal damage that occur independent of MLCK-mediated TJ breakdown. These

findings reflect the therapeutic limitations of MLCK inhibitors¹⁵⁵. Current MLCK inhibitors also lack specificity towards the catalytic domain of epithelial and smooth muscle MLCK, thus resulting in potential toxic side effects¹⁵⁶.

Potential targets for TJ regulation include occludin and claudin, both of which are important for sustaining intestinal integrity. A study on protein tyrosine phosphatase (TCPTP) activity in epithelial cells showed that TCPTP protects the intestinal barrier by directly restricting STAT1-mediated transcription of the cation-selective pore claudin-2, as well as upregulating inhibitory matriptase, which prevents occludin mislocalization and decreases epithelial permeability¹⁵⁷. However, there are currently no clinical trials testing medicines specifically targeting TJ proteins. This implies the need for further research and development in this area.

6.2. Ecological regulation of gut microbiota and the host-microbiome interaction

Probiotics have been found to regulate barrier function through multiple mechanisms. For instance, *Lactobacillus reuteri* activates the Wnt/ β -catenin signaling pathway to induce Paneth cell differentiation and AMP secretion, and meanwhile stimulates the proliferation of *Lgr5*⁺ ISCs to promote epithelial repair¹⁵⁸. Additionally, flagellin from *Roseburia* spp. binds to Toll-like receptor 5 (TLR5), which can upregulate occludin and MUC2, thereby improving intestinal barrier integrity and increasing IL-22 and REG3 γ levels to further regulate gut ecology¹⁵⁹. *Akkermansia muciniphila* also supports intestinal epithelial repair, proliferation, differentiation, and stability via the regulation of ISC program⁹⁷. Therefore, targeting intestinal microbial dysbiosis emerges as a promising therapeutic strategy for restoring impaired barrier function. Recent advancements include genetically engineered probiotics coated with overexpressed artificial enzymes using biological nanomaterials, a novel approach to mucosal repair and inflammation therapy^{160,161}. *Bifidobacterium longum*, modified to express catalase and superoxide dismutase, demonstrates enhanced colonization of the intestines, robust anti-inflammatory activity, facilitates intestinal barrier remodeling, and regulates the microbial balance. These developments could potentially reduce the adverse reactions of traditional anti-inflammatory drugs¹⁶².

BAs serve as a crucial mediator in host-microbiota communication and the regulation of gut microbiome composition¹⁶³. Dysregulation of BA synthesis is strongly associated with the pathogenesis of various diseases. Activation of FXR effectively mitigates the detrimental effects of BA overload by inhibiting cholesterol metabolism and promoting BA transportation from hepatocytes, thereby reducing cholestatic liver damage. Studies have shown that the FXR agonist obeticholic acid (OCA) can increase the proportion of microbes involved in DNA synthesis and amino acid metabolism, reduce the proliferation of certain Gram-positive bacteria, and decrease the concentrations of endogenous bile acids, thus effectively improving gut ecological disorders¹⁶⁴. OCA treatment also prevents damage to the intestinal vascular barrier by stabilizing β -catenin within endothelial cells¹⁶⁵.

AHR, a receptor for tryptophan metabolites, plays an important role in maintaining the integrity of the intestinal barrier. Studies have revealed a significant loss of epithelial barrier function in *Ahr*^{-/-} mice, characterized by destruction of the intestinal mechanical barrier and a failure to commit to a differentiating lineage¹⁶⁶. Notably, this phenotype is also observed in mice with an AHR ligand-deficient diet. Subsequent research

Table 2 Clinical trials related to intestinal barrier strengthening.

Class	Intervention	Disease	Effect on intestinal barrier function	Trial registration number
Gut ecological regulation	Rifaximin- α (550 mg BID)	Hepatic encephalopathy	↑ Faecal TNF- α , IL-17E, and IL-17A enriched intestinal microenvironment; ↓ Circulating neutrophil TLR-4 expression and plasma TNF- α ↓ Species with mucin-degrading capacity in faeces	NCT02019784
	FMT from healthy donors	DSPN	↑ ZO-1, Claudin-1, Claudin-4 in the colonic mucosa ↓ Serum LBP, IL-6, TNF- α	ChiCTR1800017257
	Non-absorbable, gut-restricted, engineered carbon bead adsorbent Yaq-001	Cirrhosis	↓ Gut permeability in the organoids	NCT03202498
Immune regulation	IL-22 Fc fusion protein UTTR1147A	UC	↑ Ducible epithelial genes <i>DMBT1</i> , <i>MUC1</i> , <i>STAT3</i> , <i>MUC4</i> , <i>REG1A</i> , and <i>REG3A</i> ↓ Baseline dysbiosis	NCT02749630
	Proline hydroxylase domain (PHD) inhibitor	IBD	Preclinical IBD models show ↓ Plasma FITC-dextran ↑ Barrier-protective genes <i>Tff3</i> , <i>Tjp1</i> , <i>Nt5e</i> ↓ proportions of monocytes, neutrophils and proinflammatory CD3 ⁺ T cell subsets; ↓ IL-17 signaling, fecal lipocalin 2 (LCN2) and calprotectin levels	CTR20241789, NCT06012578
	Transglutaminase 2 inhibitor ZED1227	Celiac disease	↓ Plasma IL-6 and TNF ↓ Ratio of villus height to crypt depth ↓ Intraepithelial lymphocyte density, Celiac Symptom Index score, and Celiac Disease Questionnaire score (100-mg dose)	EudraCT 2017-002241-30
Dietary intervention	Gluten-free diet	Irritable bowel syndrome-diarrhea	↓ Bowel movement, 0–2 h levels of mannitol and LMR; ↑ ZO-1, claudin-1, occludin in small bowel mucosa	NCT01094041
	Resistant starch intervention	Overweight or obesity	↓ MCP-1, IL-1 β , IL-6; ↑ anti-inflammatory IL-10 in serum and mWAT; ↑ TJ expression including ZO-1, Occludin; ↓ Serum DX-4000-FITC, LPS in serum and mWAT	ChiCTR-TTRCC-13003333
	Crohn's disease exclusion diet (CDED) coupled with partial enteral nutrition (PEN)	Mild to moderate Crohn's disease	↑ Proportion of patients with normal LMR; ↓ Serum level of C-reactive protein and fecal level of calprotectin; ↓ Fecal <i>Proteobacteria</i>	NCT01728870
	Oral glutamine intervention	Post-infectious diarrhea in irritable bowel syndrome	↓ LMR; ↓ Irritable Bowel Syndrome Severity Scoring (IBS-SS) scores; ↓ Daily bowel movement frequency and Bristol Stool Scale	NCT01414244

demonstrated that AHR ligands derived from diet could effectively prevent diseases that impair barrier function by upregulating the expression of TJ proteins *via* promoting intracellular Zn²⁺ signaling¹⁶⁷. Other AHR agonists, such as Urolithin A (UroA), a microbial metabolite, exhibit promise in protecting the barrier function. UroA has anti-inflammatory activity and promotes barrier repair by activating the AhR–Nrf2-dependent

pathway to upregulate TJs¹⁶⁸. These studies demonstrate the importance of intestinal AHR pathway-dependent activation in the treatment of ALD, a highly associated form of liver disease characterized by loss of TJ barriers^{169,170}. These findings suggest AHR may serve as a valuable therapeutic target for intestinal barrier repair, especially in chronic diseases associated with barrier dysfunction.




Figure 4 Potential interventions targeting the intestinal barrier to enhance the management of chronic gut and liver diseases. While immune-targeted therapies, such as anti-inflammatory agents, immunosuppressants and the biologics, can effectively reduce intestinal inflammation, they often carry the risk of serious side effects and fail to directly address barrier damage. This limitation will cause the lack of long-term cure for the disease. Therefore, it is necessary to develop effective therapeutic strategies for the timely repair of the intestinal epithelial barrier. Studies targeting TJs, ecological regulation of gut microbiota, intestinal stem cell regeneration, and immune regulation have shown potential value in enhancing enterohepatic function. Targeting intestinal epithelial TCPTP has been demonstrated to preserve intestinal barrier integrity by modulating TJ components, including normalizing claudin-2 expression and maintaining localization of ZO-1 and occludin¹⁵⁷. Probiotic administration (e.g., *Lactobacillus reuteri*, *Akkermansia muciniphila*, *Bifidobacterium longum*)^{97,158,162} and microbial metabolites^{169,170,174}, such as tryptophan derivatives, can restore intestinal barrier integrity by enhancing *Lgr5*⁺ ISCs proliferation and upregulating levels of TJ proteins, MUC2, AMPs represented by REG3γ. Emerging therapeutic strategies targeting ISCs regeneration, including modulation of the bile acid–FXR axis and organoid-based transplantation, hold significant promise for ameliorating chronic enterohepatic diseases^{72,73,180}. Additionally, immune signaling pathways mediated by cytokines such as IL-10^{183,184} and IL-22^{185,186,188} have been identified as potential therapeutic targets for enhancing intestinal barrier integrity. Created with Biorender.com.

The inhibition of IDO, an enzyme that catalyzes the conversion of tryptophan to Kyn, has shown potential for treating barrier dysfunction. By diverting tryptophan towards indole metabolism, IDO inhibition can increase the production of AHR ligands, which are beneficial for intestinal integrity. While IDO inhibitors demonstrated favorable outcomes in cancer therapy¹⁷¹, their application in managing intestinal barrier damage requires further investigation. The IDO inhibitor indoximod has been shown to effectively preserve tight intercellular connections and significantly improve DSS-induced colon injury¹⁷². However, it is important to consider that IDO inhibition might also induce adverse enterotoxic effects, suggesting that complete blockade of the Kyn pathway may, paradoxically, compromise epithelial barrier integrity. Studies have shown that chemotherapeutics-induced intestinal damage triggers an upregulation of tryptophan–Kyn–kynurenic acid pathway, which promotes the production of kynurenic acid in gut, and subsequently feeds back to

activate GPR35 to enhance intestinal integrity¹⁷³. These findings imply that indiscriminate inhibition of Kyn pathway could dampen this GPR35-mediated protective response by reducing kynurenic acid levels, and potentially decelerate the barrier repair. Such complexities of tryptophan metabolism in gut health suggest the need for comprehensive research when developing new therapeutic strategies targeting intestinal barrier dysfunction.

While substantial research has demonstrated that direct modulation of gut microbiota improves intestinal barrier function and ameliorates enterohepatic diseases, microbial-based therapies remain limited in clinical practice. The intestinal barrier-enhancing properties of microbiome-derived beneficial metabolites, such as indole-3-lactic acid (ILA), are contingent upon host-specific microbiome regulation¹⁷⁴. Clinical efficacy is fundamentally determined by baseline host microbiota composition, which governs both microbial colonization dynamics¹⁷⁵. Furthermore, host genetic variation constitutes a critical determinant of

therapeutic outcomes. For instance, IBD patients carrying CARD9 risk alleles exhibit impaired microbial conversion of dietary tryptophan into AHR-activating metabolites¹⁷⁶. These findings elucidate the molecular basis for the substantial interindividual variability observed in microbiota-targeted therapies. Probiotics, as symbiotic microorganisms depleted during disease states, require a stable host microenvironment for colonization and function. Without prior restoration of the impaired gut ecosystem, microbial intervention therapies often fail to establish durable microbial equilibrium or achieve meaningful clinical outcomes¹⁷⁷. Consequently, future therapeutic strategies should adopt a dual-target approach combining microbial modulation with microenvironment restoration, designed to synergistically re-establish gut ecological balance and promote epithelial barrier healing.

6.3. Intestinal stem cell regeneration

The renewal, repair, and regeneration of intestinal epithelium are heavily dependent on *Lgr5*⁺ ISC, located at the crypt base. *Lgr5*, a specific marker for ISCs, encodes receptors that can trigger the reprogramming of ISCs into epithelial lineages in response to signals such as Wnt¹⁷⁸. ISC activity, regulated by both extracellular and paracrine signals, maintains homeostasis and initiates adaptive differentiation in response to injury to ensure the fundamental physiological functions of the intestine. BAs play a crucial role in mediating gut–liver communication and integrate dietary and metabolic signals that regulate ISC function⁷³. Recent studies have shown a novel intervention for treating IBD-associated intestinal damage by reducing excessive BA accumulation in gut under pathological conditions. FXR agonist treatment reduces the enterohepatic levels of cholic acid by suppressing CYP8B1 expression in the liver, and thus alleviates the BA-related toxicity on *Lgr5*⁺ ISCs⁷². These findings suggest the feasibility of targeting ISCs to treat IBD based on cross-organ metabolic regulation.

The significant role of ISCs in injury repair and self-renewal has also led to the exploration of ISC culture and transplantation to the site of injury as a therapeutic strategy for repairing damaged mucosa. For instance, colon organoids derived from *Lgr5*⁺ ISCs successfully restore damaged rectal epithelium through enema transplantation¹⁷⁹. Moreover, transplantation of ileum-derived organoids into colons has shown potential in creating functional small intestinalized colon (SIC) tissue, which retains regional characteristics such as absorptive functions, presenting promise for the treatment of short bowel syndrome (SBS)¹⁸⁰. The therapeutic applications of organoids, with their ease of *in vitro* culture and physiological resemblance to native tissue, have opened new avenues in regenerative medicine. However, there are many translation challenges that need to be addressed, such as efficient *in vivo* delivery, potential immune rejection, and the long-term stability of transplanted cells. These issues warrant further investigation. The potential of intestinal organoids in regenerative medicine is vast, and continued research in this field might lead to substantial breakthroughs in the treatment of gastrointestinal and extra-intestinal diseases.

6.4. Immune regulation

Traditional anti-inflammatory therapies, including corticosteroids, 5-aminosalicylates, and the novel TNF- α monoclonal antibodies, have been the standard treatments for IBD and other immune-mediated disorders. While these therapies effectively reduce

inflammation and alleviate symptoms, they often fall short in achieving long-term healing, particularly in relation to the intestinal barrier. Many patients experience a loss of response over time and disease recurrence. For instance, although anti-IL-6 antibody therapy was effective in moderate to severe CD or UC in clinical trials, serious adverse reactions such as abscesses and intestinal perforations continue to occur in some patients¹⁸¹. These therapies often fail to address the underlying epithelial dysfunction, which is a key factor in sustaining chronic intestinal inflammation.

To address these limitations, there is growing interest in targeting immune pathways that are more directly involved in epithelial barrier repair. IL-10 is an effective anti-inflammatory and tissue-regenerative cytokine¹⁸², and recent IL-10 preparations have been developed with the goal of enhanced therapeutic effects by targeting mucosal barriers^{183,184}. Another promising target is IL-22, which has its receptor IL-22R expressed on intestinal epithelial cells, making IL-22 and IL-22R potential candidates for therapies aimed at restoring epithelial integrity and enhancing mucosal healing. IL-22 plays a pivotal role in maintaining intestinal homeostasis by stimulating the production of AMPs and mucins, as well as ISC regeneration. Consequently, IL-22 is considered a potential therapeutic approach for barrier repair. IL-22 controls the generation of Paneth cell generation, thereby increasing intestinal levels of AMPs¹⁸⁵. Moreover, IL-22 initiates the transcription of IL-18 *via* STAT3 phosphorylation, which promotes the proliferation and repair capability of *Lgr5*⁺ ISCs and the bacterial clearance by IFN- γ ⁺ T cells *via* Akt–Tcf4 signaling pathway to enhance the epithelial barrier defense¹⁸⁶.

The IL-22 fusion protein agonist Efmardocokin alfa (UTTR1147A) is currently under investigation for active UC and CD treatment (NCT02749630). Clinical trials have shown that UTTR1147A activates IL-22R signaling pathways in both UC patients and healthy individuals, with improvements in UC-associated dysbiosis¹⁸⁷. Furthermore, specific activation of IL-22 signaling in the gut can enhance hepatic and systemic glucose and lipid metabolic homeostasis in a microbiota-dependent manner in models of metabolic disorders¹⁸⁸. Additionally, IL-22 has also demonstrated positive effects on MASLD, ALD and dietary obesity^{159,189}. When administered exogenously, IL-22 exerts its therapeutic effects through its receptor on IECs, rather than hepatocytes, and then activates STAT3 and inhibits WNT– β -catenin signaling to reduce the absorptive enterocyte compartment. Nonetheless, the role of IL-22 in intestinal barrier maintenance remains controversial, with concerns over possible pathogenic immunomodulatory effects, such as mediating CXCR2⁺ neutrophil chemotaxis in colonic tissue and increasing resistance to the IL-23 monoclonal antibody Ustekinumab. These observations indicate that IL-22-targeted therapies might not always achieve the desired efficacy, necessitating further research to better understand the full impact and potential side effects of IL-22 activation.

IL-17A, mainly produced by Th17 cells, mediates inflammatory infiltration and the recruitment of immune cells. Although anti-IL-17A monoclonal antibodies have been used to treat autoimmune diseases, their application in IBD has led to significant adverse reactions and, in some cases, disease exacerbation. Research has revealed the essential role of IL-17A signaling in the expression of ATOH1, the key transcription factor that regulates the differentiation of *Lgr5*⁺ ISCs into secretory cell lineages vital for maintaining mucosal epithelial integrity¹⁹⁰. This observation highlights the challenges of relying solely on immunotherapy, as

blocking IL-17A signaling, can disrupt epithelial homeostasis and further induce drug resistance and disease recurrence. Additionally, potential immune targets for promoting mucosal healing include CXCL1 production mediated by intestinal PTGER4⁺ macrophages, which has been associated with enhanced tissue repair. Furthermore, macrophage-derived TGFB1 has been shown to activate pro-regenerative transcription factors such as YAP/TEAD and SOX9, which promote epithelial regeneration *via* signaling pathways critical for ISC function¹⁹¹. Therefore, a deeper understanding of intercellular communication mechanisms, particularly among immune cells, ISCs, and intestinal secretory cells, will be necessary in the future. The approach, which combines the synergistic actions of immune and regenerative signals, holds great promise for the development of next-generation treatments for IBD and other gut and liver disorders.

6.5. Potential advances in clinical trials related to intestinal barrier function strengthening

Multiple innovative therapies targeting intestinal barrier integrity are currently undergoing clinical validation (Table 2). The novel gut-restricted selective prolyl hydroxylase domain (PHD) inhibitor ISM5411, developed using a multimodal generative artificial intelligence platform, has successfully completed Phase I trials (CTR20241789, NCT06012578). Preclinical studies demonstrate the dual mechanism of this drug in intestinal mucosal repair and immune regulation, showing significant efficacy in IBD models¹⁹². Global multicenter trials are now planned to further validate these therapeutic benefits. Clinical evidence demonstrates that Rifaximin- α effectively enhances intestinal barrier repair by modulating the gut microenvironment through upregulation of TNF- α and interleukin-17E in feces, thereby strengthening antimicrobial defenses against pathogenic invasion (NCT02019784)¹¹². Furthermore, fecal microbiota transplantation (FMT) from healthy donors has shown therapeutic potential in restoring intestinal barrier function and mitigating systemic inflammation in diabetic distal symmetric polyneuropathy (DSPN)¹⁹³. A non-absorbable, gut-restricted, engineered carbon bead adsorbent Yaq-001 exhibits clinically relevant efficacy in cirrhosis by ameliorating gut barrier dysfunction and reducing systemic endotoxin burden (NCT03202498)¹⁹⁴. Dietary interventions for barrier repair have demonstrated efficacy across multiple gastrointestinal conditions. Clinical studies show benefits in irritable bowel syndrome-diarrhea (NCT01094041)¹⁹⁵, metabolic disorders (ChiCTR-TTRCC-13003333)¹⁹⁶, and mild to moderate Crohn's disease (NCT01728870)¹⁹⁷. Notably, dietary glutamine supplementation significantly restored intestinal permeability and alleviated post-infectious diarrhea in irritable bowel syndrome (NCT01414244)¹⁹⁸. In celiac disease, the transglutaminase 2 inhibitor ZED1227 demonstrated significant improvement in duodenal mucosal architecture, including villous height and crypt depth, while reducing intraepithelial lymphocyte infiltration. The therapeutic mechanism involves inhibition of glutamine residue deamidation in immunogenic gluten peptides, thereby preventing the activation of T cell and subsequent mucosal injury (EudraCT 2017-002241-30)^{199,200}. Engineered probiotics demonstrate significant therapeutic potential for restoring intestinal barrier integrity and maintaining mucosal homeostasis. Current research focuses on combining their barrier-enhancing properties with immunomodulatory effects, though clinical translation remains at the preclinical stage. While these advances underscore the promise of targeted barrier repair

strategies, comprehensive evaluation of long-term safety and efficacy in human trials is needed to advance clinical applications.

7. Conclusions and perspectives

The intestinal barrier is a complex and dynamic structure that maintains gut homeostasis, mediates host-microbiome interactions, and shields against pathogens and toxins. Its dysfunction is central to the development of many gastrointestinal diseases, including IBD, and extra-intestinal conditions such as obesity and MASLD. Emerging research on gut–X communication emphasizes the need for therapeutic strategies that not only reduce inflammation but also promote epithelial barrier repair. While advances in immunosuppressive and biologic therapies for IBD and related disorders have improved symptom management, they often fail to address the root cause of barrier dysfunction. Prolonged use of these treatments can lead to significant side effects, including infection risks, malignancy, and disease recurrence, suggesting the need for a paradigm shift toward therapies that integrate barrier-restorative strategies with traditional anti-inflammatory approaches. Stem cell-based regeneration, microbial therapy, bile acid modulation, and TJ modulators are promising for achieving durable disease remission.

Intestinal barrier dysfunction exhibits a complex bidirectional causality with disease pathogenesis, serving as a primary trigger initiating pathological cascades or as a consequence exacerbating disease progression. This dynamic interplay is critically influenced by genetic susceptibility, environmental exposures, and stage-specific diseases. As a pathogenic cause, intestinal barrier dysfunction precedes overt pathological manifestations and initiates disease processes through diverse mechanisms mediated by genetic predisposition, dietary factors, circadian rhythm, drug usage, and other environmental influences. Disruption of the mucus layer and TJ structures enables pathogenic bacteria and their metabolites to translocate into systemic circulation, triggering inflammatory response and multi-organ dysfunction that ultimately contribute to the pathogenesis of MASH, metabolic disorders, IBD, CRC, ALD, and PSC. Intestinal barrier injury is also a pathological consequence and disease amplifier, generating multi-organ pathological cascades through gut dysbiosis, immune hyperactivation, and metabolite toxicity. These mechanisms exacerbate conditions including IBD, MASH, PSC, and CRC. Furthermore, the resulting disease microenvironment reciprocally impairs barrier integrity, establishing a self-perpetuating vicious cycle of pathophysiology.

There is accumulating clinical evidence of off-target and systemic side effects from immunosuppressive therapies, including the loss of treatment efficacy, exacerbated inflammatory damage, and immune disorders²⁰¹. Inflammatory cytokines can directly damage the epithelial barrier to aggravate inflammation, and necessitate anti-inflammatory interventions. However, these same cytokines also trigger essential regenerative signals that promote mucosal healing²⁰². Immunosuppressive therapies may block these beneficial signals, and thereby inhibit the barrier repair process, worsen microbial imbalances, and further damage epithelial integrity. This interplay between therapeutic benefits and adverse effects complicates disease management, leading to clinical recurrence and treatment failure. While our understanding of epithelial barrier structure and function has advanced, the etiology of barrier damage and the complex network mechanisms

involved in intestinal mucosal healing remain unclear. The further exploration into intercellular communication among intestinal cells, immune, non-immune, host, and microbial, will be crucial for identifying more effective targets.

The transition from NAFLD to MASLD identified the significant role of global metabolic dysfunction in the pathogenesis of fatty liver diseases^{203,204}. Increasing evidence suggests that the loss of intestinal barrier homeostasis, which is important for systemic metabolic balance, is closely linked to liver diseases²⁰⁵. Consequently, repairing the intestinal barrier is now recognized as integral to treating liver disease, and the gut–liver axis has emerged as a promising therapeutic target. This has led to the novel concept of “enteric treatment for liver disease”, with the BA-intestinal FXR signaling pathway demonstrating considerable efficacy in MASLD treatment^{206,207}. Furthermore, there are still many unknown enterohepatic communication secretory proteins and metabolites that warrant further exploration.

In addition to pharmacological treatments, regenerative medicine has progressed rapidly with the development of stem cell-derived organoids. These organoids, which can be cultured and transplanted to mimic the functional properties of *in vivo* tissue, offer new possibilities for repairing damaged gut epithelium. They hold particular promise for conditions where endogenous regeneration is insufficient, such as refractory IBD or SBS. However, challenges remain in the therapeutic application of organoids, such as ensuring functional integration, preventing immune rejection, and achieving long-term stability post-transplantation. Looking ahead, a comprehensive approach combining ISC-targeted therapies, microbial modulation, immune regulation, and organoid-based regeneration offers a multifaceted pathway to improve outcomes in diseases linked to intestinal barrier dysfunction. Future research should focus on developing precise interventions that work synergistically with existing treatments to promote barrier repair, restore homeostasis, and prevent relapse.

When translating experimental data to human disease pathogenesis, it is crucial to carefully evaluate the animal models used. The DSS colitis model specifically induces damage to the intestinal epithelial barrier through TJ dependence, but it fails to reflect the damage observed in non-TJ dependent regions of the human intestinal tract in conditions as IBD. Meanwhile, the types and abundance of barrier injury markers used in animal models must align with the clinical characteristics of patients to ensure the translational relevance of experimental findings.

Barrier defense and repair, independent of immunosuppression, is a valuable objective for treating hepatenteric diseases. Specific intestinal barrier disorders vary across diseases and individuals, suggesting that personalized treatment strategies may be more effective. Enhancing intestinal barrier integrity has been shown to have great potential in preventing and reversing liver steatosis, immune infiltration and fibrosis in MASLD²⁰⁸. For milder conditions, improvements and prevention can be achieved through dietary or non-pharmacological methods, such as probiotics and related products. In the future, human studies will be critical to validate these potential targets and treatments (Table 1)^{37,64–70,72,73,89,95–97,99,106,108,113,118,133,134,136–138,140,146–148,151–153,209}.

In conclusion, restoring the intestinal barrier represents is not only a therapeutic challenge, but also a significant opportunity. By addressing the fundamental defects in epithelial integrity and the complex interactions that maintain it, we stand on the verge of a paradigm shift in the treatment of intestinal and systemic diseases.

The convergence of novel biotechnologies, regenerative medicine, and microbiome research promises to reshape the landscape of gastrointestinal therapy and holds great potential to improve patient outcomes.

Acknowledgements

This study was supported by National Natural Science Foundation of China (82530024 and 82222071), National Key Research and Development Program of China (2021YFA1301200), Strategic Priority Research Program of the Chinese Academy of Sciences (XDB0830402, China), and Shanghai Municipal Science and Technology Major Project, to Cen Xie.

Author contributions

Yongxin Zhang: Writing – review, editing, Writing – original draft, Visualization, Methodology, Conceptualization. Yameng Liu: Writing – review, editing, Project administration, Conceptualization. Xinyu Liang: Writing – review, editing. Yingquan Wen: Writing – review, editing. Jingjie Zhao: Writing – review, editing. Yong He: Writing – review, editing, Conceptualization. Qing Xie: Writing – review, editing, Conceptualization. Cen Xie: Writing – review, editing, Supervision, Project administration, Funding acquisition, Conceptualization.

Conflicts of interest

The authors declare no conflicts of interest.

References

- Leibovitz H, Lee SH, Raygoza Garay JA, Espin Garcia O, Xue MY, Neustaeter A, et al. Immune response and barrier dysfunction-related proteomic signatures in preclinical phase of Crohn's disease highlight earliest events of pathogenesis. *Gut* 2023;**72**:1462–71.
- Shin W, Kim HJ. Intestinal barrier dysfunction orchestrates the onset of inflammatory host-microbiome cross-talk in a human gut inflammation-on-a-chip. *Proc Natl Acad Sci U S A* 2018;**115**:E10539–47.
- Vanheel H, Vicario M, Vanuytsel T, Van Oudenhove L, Martinez C, Keita ÁV, et al. Impaired duodenal mucosal integrity and low-grade inflammation in functional dyspepsia. *Gut* 2014;**63**:262–71.
- Kiesslich R, Duckworth CA, Moussata D, Gloeckner A, Lim LG, Goetz M, et al. Local barrier dysfunction identified by confocal laser endomicroscopy predicts relapse in inflammatory bowel disease. *Gut* 2012;**61**:1146–53.
- Sözener ZC, Cevhertas L, Nadeau K, Akdis M, Akdis CA. Environmental factors in epithelial barrier dysfunction. *J Allergy Clin Immunol* 2020;**145**:1517–28.
- Lopez Otin C, Kroemer G. Hallmarks of health. *Cell* 2021;**184**:33–63.
- Swidsinski A, Loening-Baucke V, Theissig F, Engelhardt H, Bengmark S, Koch S, et al. Comparative study of the intestinal mucus barrier in normal and inflamed colon. *Gut* 2007;**56**:343–50.
- van der Post S, Jabbar KS, Birchenough G, Arike L, Akhtar N, Sjövall H, et al. Structural weakening of the colonic mucus barrier is an early event in ulcerative colitis pathogenesis. *Gut* 2019;**68**:2142–51.
- Johansson MEV, Phillipson M, Petersson J, Velcich A, Holm L, Hansson GC. The inner of the two muc2 mucin-dependent mucus layers in colon is devoid of bacteria. *Proc Natl Acad Sci U S A* 2008;**105**:15064–9.
- Tsukita S, Furuse M, Itoh M. Multifunctional strands in tight junctions. *Nat Rev Mol Cell Biol* 2001;**2**:285–93.

11. Metidji A, Omenetti S, Crotta S, Li Y, Nye E, Ross E, et al. The environmental sensor AHR protects from inflammatory damage by maintaining intestinal stem cell homeostasis and barrier integrity. *Immunity* 2019;**50**:1542.
12. Kinchen J, Chen HH, Parikh K, Antanaviciute A, Jagielowicz M, Fawcner Corbett D, et al. Structural remodeling of the human colonic mesenchyme in inflammatory bowel disease. *Cell* 2018;**175**:372–86.e17.
13. Spalinger MR, Sayoc-Becerra A, Santos AN, Shawki A, Canale V, Krishnan M, et al. Ptpn2 regulates interactions between macrophages and intestinal epithelial cells to promote intestinal barrier function. *Gastroenterology* 2020;**159**:1763–77.e14.
14. Kayama H, Okumura R, Takeda K. Interaction between the microbiota, epithelia, and immune cells in the intestine. *Annu Rev Immunol* 2020;**38**:23–48.
15. Rouland M, Beaudoin L, Rouxel O, Bertrand L, Cagninacci L, Saffarian A, et al. Gut mucosa alterations and loss of segmented filamentous bacteria in type 1 diabetes are associated with inflammation rather than hyperglycaemia. *Gut* 2022;**71**:296–308.
16. Leonardi I, Gao IH, Lin WY, Allen M, Li XV, Fiers WD, et al. Mucosal fungi promote gut barrier function and social behavior via type 17 immunity. *Cell* 2022;**185**:831–46.e14.
17. Guin J, Hannezo E, Yui SR, Denharther S, Ulyanchenko S, Maimets M, et al. Tracing the origin of adult intestinal stem cells. *Nature* 2019;**570**:107–11.
18. Haber AL, Biton M, Rogel N, Herbst RH, Shekhar K, Smillie C, et al. A single-cell survey of the small intestinal epithelium. *Nature* 2017;**551**:333–9.
19. Nystrom EEL, Martinez Abad B, Arike L, Birchenough GMH, Nonnecke EB, Castillo PA, et al. An intercrypt subpopulation of goblet cells is essential for colonic mucus barrier function. *Science* 2021;**372**:eabb1590.
20. Gerbe F, Sidot E, Smyth DJ, Ohmoto M, Matsumoto I, Dardalhon V, et al. Intestinal epithelial tuft cells initiate type 2 mucosal immunity to helminth parasites. *Nature* 2016;**529**:226–30.
21. Wolf JL, Rubin DH, Finberg R, Kauffman RS, Sharpe AH, Trier JS, et al. Intestinal m cells: a pathway for entry of reovirus into the host. *Science* 1981;**212**:471–2.
22. Besselink MG, van Santvoort HC, Renooij W, de Smet MB, Boermeester MA, Fischer K, et al. Intestinal barrier dysfunction in a randomized trial of a specific probiotic composition in acute pancreatitis. *Ann Surg* 2009;**250**:712–9.
23. Linh HT, Iwata Y, Senda Y, Sakai Takemori Y, Nakade Y, Oshima M, et al. Intestinal bacterial translocation contributes to diabetic kidney disease. *J Am Soc Nephrol* 2022;**33**:1105–19.
24. Reyes H, Zapata R, Hernández I, Gotteland M, Sandoval L, Jirón MI, et al. Is a leaky gut involved in the pathogenesis of intrahepatic cholestasis of pregnancy?. *Hepatology* 2006;**43**:715–22.
25. Watson AJM. Claudins and barrier dysfunction in intestinal inflammation: cause or consequence?. *Gut* 2015;**64**:1501–2.
26. Schreiber S, Aden K, Bernardes JP, Conrad C, Tran F, Höper H, et al. Therapeutic interleukin-6 trans-signaling inhibition by olamicept (sgp130fc) in patients with active inflammatory bowel disease. *Gastroenterology* 2021;**160**:2354–66.e11.
27. Olivera PA, Lasa JS, Bonovas S, Danese S, Peyrin-Biroulet L. Safety of janus kinase inhibitors in patients with inflammatory bowel diseases or other immune-mediated diseases: a systematic review and meta-analysis. *Gastroenterology* 2020;**158**:1554–73.e12.
28. Sommer K, Heidbreder K, Kreiss L, Dedden M, Paap EM, Wiendl M, et al. Anti- $\beta 7$ integrin treatment impedes the recruitment on non-classical monocytes to the gut and delays macrophage-mediated intestinal wound healing. *Clin Transl Med* 2023;**13**:e1233.
29. Horowitz A, Chanez-Paredes SD, Haest X, Turner JR. Paracellular permeability and tight junction regulation in gut health and disease. *Nat Rev Gastroenterol Hepatol* 2023;**20**:417–32.
30. Gustafsson JK, Johansson MEV. The role of goblet cells and mucus in intestinal homeostasis. *Nat Rev Gastroenterol Hepatol* 2022;**19**:785–803.
31. Peterson LW, Artis D. Intestinal epithelial cells: regulators of barrier function and immune homeostasis. *Nat Rev Immunol* 2014;**14**:141–53.
32. Paone P, Cani PD. Mucus barrier, mucins and gut microbiota: the expected slimy partners?. *Gut* 2020;**69**:2232–43.
33. Schutte A, Ermund A, Becker-Pauly C, Johansson ME, Rodriguez-Pineiro AM, Backhed F, et al. Microbial-induced meprin beta cleavage in muc2 mucin and a functional cfr channel are required to release anchored small intestinal mucus. *Proc Natl Acad Sci U S A* 2014;**111**:12396–401.
34. Luis AS, Hansson GC. Intestinal mucus and their glycans: a habitat for thriving microbiota. *Cell Host Microbe* 2023;**31**:1087–100.
35. Katoh T, Yamada C, Wallace MD, Yoshida A, Gotoh A, Arai M, et al. A bacterial sulfoglycosidase highlights mucin O-glycan breakdown in the gut ecosystem. *Nat Chem Biol* 2023;**19**:778–89.
36. Bergstrom K, Shan X, Casero D, Batushansky A, Lagishetty V, Jacobs JP, et al. Proximal colon-derived O-glycosylated mucus encapsulates and modulates the microbiota. *Science* 2020;**370**:467–72.
37. Desai MS, Seekatz AM, Koropatkin NM, Kamada N, Hickey CA, Wolter M, et al. A dietary fiber-deprived gut microbiota degrades the colonic mucus barrier and enhances pathogen susceptibility. *Cell* 2016;**167**:1339–53.e21.
38. Kissoon-Singh V, Moreau F, Trusevych E, Chadee K. Entamoeba histolytica exacerbates epithelial tight junction permeability and proinflammatory responses in muc2^{-/-} mice. *Am J Pathol* 2013;**182**:852–65.
39. Zhang JH, Huang Y, Li H, Xu PF, Liu QH, Sun Y, et al. B3galt5 functions as a pxx target gene and regulates obesity and insulin resistance by maintaining intestinal integrity. *Nat Commun* 2024;**15**:5919.
40. Zhang J, Yu QZ, Jiang DL, Yu K, Yu WW, Chi ZX, et al. Epithelial gasdermin d shapes the host-microbial interface by driving mucus layer formation. *Sci Immunol* 2022;**7**:eabk2092.
41. Liang W, Enee E, Andre Vallee C, Falcone M, Sun J, Diana J. Intestinal cathelicidin antimicrobial peptide shapes a protective neonatal gut microbiota against pancreatic autoimmunity. *Gastroenterology* 2022;**162**:1288–302.e16.
42. Belkaid Y, Hand TW. Role of the microbiota in immunity and inflammation. *Cell* 2014;**157**:121–41.
43. Martinez Lopez M, Iborra S, Conde Garrosa R, Mastrangelo A, Danne C, Mann ER, et al. Microbiota sensing by mincle—syk axis in dendritic cells regulates interleukin-17 and -22 production and promotes intestinal barrier integrity. *Immunity* 2019;**50**:446–61.e9.
44. Stojanovic O, Miguel-Aliaga I, Trajkovski M. Intestinal plasticity and metabolism as regulators of organismal energy homeostasis. *Nat Metab* 2022;**4**:1444–58.
45. VanDussen KL, Stojmirovic A, Li K, Liu TC, Kimes PK, Muegge BD, et al. Abnormal small intestinal epithelial microvilli in patients with Crohn's disease. *Gastroenterology* 2018;**155**:815–28.
46. Huang CR, Kuo CJ, Huang CW, Chen YT, Liu BY, Lee CT, et al. Host CDK-1 and formin mediate microvillar effacement induced by enterohemorrhagic escherichia coli. *Nat Commun* 2021;**12**:90.
47. Otani T, Furuse M. Tight junction structure and function revisited: (trends in cell biology 30, 805-817, 2020). *Trends Cell Biol* 2020;**30**:1014.
48. Zihni C, Mills C, Matter K, Balda MS. Tight junctions: from simple barriers to multifunctional molecular gates. *Nat Rev Mol Cell Biol* 2016;**17**:564–80.
49. Tsukita S, Tanaka H, Tamura A. The claudins: from tight junctions to biological systems. *Trends Biochem Sci* 2019;**44**:141–52.
50. Raju P, Shashikanth N, Tsai PY, Pongkorpsakol P, Chanez Paredes S, Steinhagen PR, et al. Inactivation of paracellular cation-selective claudin-2 channels attenuates immune-mediated experimental colitis in mice. *J Clin Invest* 2020;**130**:5197–208.
51. Zeisel MB, Dhawan P, Baumert TF. Tight junction proteins in gastrointestinal and liver disease. *Gut* 2019;**68**:547–61.
52. Kuo WT, Zuo L, Odenwald MA, Madha S, Singh G, Gurniak CB, et al. The tight junction protein ZO-1 is dispensable for barrier

- function but critical for effective mucosal repair. *Gastroenterology* 2021;**161**:1924–39.
53. Holthaus D, Kraft MR, Krug SM, Wolf S, Muller A, Delgado Betancourt E, et al. Dissection of barrier dysfunction in organoid-derived human intestinal epithelia induced by giardia duodenalis. *Gastroenterology* 2022;**162**:844–58.
 54. Dolat L, Carpenter VK, Chen YS, Suzuki M, Smith EP, Kuddar O, et al. Chlamydia repurposes the actin-binding protein eps8 to disassemble epithelial tight junctions and promote infection. *Cell Host Microbe* 2022;**30**:1685–700.e10.
 55. Kang S, Kim J, Park A, Koh M, Shin W, Park G, et al. Trim40 is a pathogenic driver of inflammatory bowel disease subverting intestinal barrier integrity. *Nat Commun* 2023;**14**:700.
 56. Gehart H, Clevers H. Tales from the crypt: new insights into intestinal stem cells. *Nat Rev Gastroenterol Hepatol* 2019;**16**:19–34.
 57. Cox CB, Storm EE, Kapoor VN, Chavarria Smith J, Lin DL, Wang LF, et al. IL-1R1-dependent signaling coordinates epithelial regeneration in response to intestinal damage. *Sci Immunol* 2021;**6**:eabe8856.
 58. Wei W, Liu Y, Hou Y, Cao S, Chen Z, Zhang Y, et al. Psychological stress-induced microbial metabolite indole-3-acetate disrupts intestinal cell lineage commitment. *Cell Metab* 2024;**36**:466–83.e7.
 59. Nalapareddy K, Nattamai KJ, Kumar RS, Karns R, Wikenheiser-Brokamp KA, Sampson LL, et al. Canonical Wnt signaling ameliorates aging of intestinal stem cells. *Cell Rep* 2017;**18**:2608–21.
 60. Gerbe F, Sidot E, Smyth DJ, Ohmoto M, Matsumoto I, Dardalhon V, et al. Intestinal epithelial tuft cells initiate type 2 mucosal immunity to helminth parasites. *Nature* 2016;**529**:226–60.
 61. Xiong Z, Zhu X, Geng J, Xu Y, Wu R, Li C, et al. Intestinal tuft-2 cells exert antimicrobial immunity via sensing bacterial metabolite *n*-undecanoylglycine. *Immunity* 2022;**55**:686–700.e7.
 62. Huang L, Bernink JH, Giladi A, Krueger D, van Son GJF, Geurts MH, et al. Tuft cells act as regenerative stem cells in the human intestine. *Nature* 2024;**634**:929–35.
 63. Tam V, Patel N, Turcotte M, Bosse Y, Pare G, Meyre D. Benefits and limitations of genome-wide association studies. *Nat Rev Genet* 2019;**20**:467–84.
 64. Sazonovs A, Stevens CR, Venkataraman GR, Yuan K, Avila B, Abreu MT, et al. Large-scale sequencing identifies multiple genes and rare variants associated with Crohn's disease susceptibility. *Nat Genet* 2022;**54**:1275–83.
 65. Gettler K, Levantovsky R, Moscatti A, Giri M, Wu Y, Hsu NY, et al. Common and rare variant prediction and penetrance of ibd in a large, multi-ethnic, health system-based biobank cohort. *Gastroenterology* 2021;**160**:1546–57.
 66. Glocker EO, Kotlarz D, Boztug K, Gertz EM, Schaffer AA, Noyan F, et al. Inflammatory bowel disease and mutations affecting the interleukin-10 receptor. *N Engl J Med* 2009;**361**:2033–45.
 67. Shao XX, Lin DP, Sun L, Wu CQ, Yang W, Jiang Y. Association of ulcerative colitis with solute-linked carrier family 26 member a3 gene polymorphisms and its expression in colonic tissues in chinese patients. *Int J Colorectal Dis* 2018;**33**:1169–72.
 68. Vinayaga-Pavan M, Frampton M, Pontikos N, Levine AP, Smith PJ, Jonasson JG, et al. Elevation in cell cycle and protein metabolism gene transcription in inactive colonic tissue from icelandic patients with ulcerative colitis. *Inflamm Bowel Dis* 2019;**25**:317–27.
 69. Kumar A, Priyamvada S, Ge Y, Jayawardena D, Singhal M, Anbazhagan AN, et al. A novel role of slc26a3 in the maintenance of intestinal epithelial barrier integrity. *Gastroenterology* 2021;**160**:1240–55.e3.
 70. Yao YK, Kim G, Shafer S, Chen ZJ, Kubo S, Ji YL, et al. Mucus sialylation determines intestinal host-commensal homeostasis. *Cell* 2022;**185**:1172–88.e28.
 71. Fazio A, Bordoni D, Kuiper JWP, Weber-Stiehl S, Stengel ST, Arnold P, et al. DNA methyltransferase 3a controls intestinal epithelial barrier function and regeneration in the colon. *Nat Commun* 2022;**13**:6266.
 72. Chen L, Jiao T, Liu W, Luo Y, Wang J, Guo X, et al. Hepatic cytochrome p450 8B₁ and cholic acid potentiate intestinal epithelial injury in colitis by suppressing intestinal stem cell renewal. *Cell Stem Cell* 2022;**29**:1366–81.e9.
 73. Sorrentino G, Perino A, Yildiz E, El Alam G, Bou Sleiman M, Gioiello A, et al. Bile acids signal via tgr5 to activate intestinal stem cells and epithelial regeneration. *Gastroenterology* 2020;**159**:956–68.e8.
 74. Langmann T, Moehle C, Mauerer R, Scharl M, Liebisch G, Zahn A, et al. Loss of detoxification in inflammatory bowel disease: dysregulation of pregnane X receptor target genes. *Gastroenterology* 2004;**127**:26–40.
 75. Venkatesh M, Mukherjee S, Wang H, Li H, Sun K, Benecet AP, et al. Symbiotic bacterial metabolites regulate gastrointestinal barrier function via the xenobiotic sensor pxxr and toll-like receptor 4. *Immunity* 2014;**41**:296–310.
 76. Huang K, Mukherjee S, DesMarais V, Albanese JM, Rafti E, Draghi Li A, et al. Targeting the pxxr-tlr4 signaling pathway to reduce intestinal inflammation in an experimental model of necrotizing enterocolitis. *Pediatr Res* 2018;**83**:1031–40.
 77. Zhou C, Tabb MM, Nelson EL, Grun F, Verma S, Sadatrafiei A, et al. Mutual repression between steroid and xenobiotic receptor and NF- κ B signaling pathways links xenobiotic metabolism and inflammation. *J Clin Invest* 2006;**116**:2280–9.
 78. Mencarelli A, Renga B, Palladino G, Claudio D, Ricci P, Distrutti E, et al. Inhibition of NF- κ B by a PXR-dependent pathway mediates counter-regulatory activities of rifaximin on innate immunity in intestinal epithelial cells. *Eur J Pharmacol* 2011;**668**:317–24.
 79. Garg A, Zhao A, Erickson SL, Mukherjee S, Lau AJ, Alston L, et al. Pregnane X receptor activation attenuates inflammation-associated intestinal epithelial barrier dysfunction by inhibiting cytokine-induced myosin light-chain kinase expression and c-jun N-terminal kinase 1/2 activation. *J Pharmacol Exp Therapeut* 2016;**359**:91–101.
 80. Makishima M, Lu TT, Xie W, Whitfield GK, Domoto H, Evans RM, et al. Vitamin D receptor as an intestinal bile acid sensor. *Science* 2002;**296**:1313–6.
 81. Kellermann L, Hansen SL, Maciag G, Granau AM, Johansen JV, Teves JM, et al. Influence of vitamin d receptor signalling and vitamin d on colonic epithelial cell fate decisions in ulcerative colitis. *J Crohns Colitis* 2024;**18**:1672–89.
 82. Chatterjee I, Zhang Y, Zhang J, Lu R, Xia Y, Sun J. Overexpression of vitamin D receptor in intestinal epithelia protects against colitis via upregulating tight junction protein claudin 15. *J Crohns Colitis* 2021;**15**:1720–36.
 83. Zhang Y, Zhang J, Xia Y, Sun J. Bacterial translocation and barrier dysfunction enhance colonic tumorigenesis. *Neoplasia* 2023;**35**:100847.
 84. Lu R, Zhang YG, Xia Y, Zhang J, Kaser A, Blumberg R, et al. Paneth cell alertness to pathogens maintained by vitamin d receptors. *Gastroenterology* 2021;**160**:1269–83.
 85. Garcia-Martinez JM, Chocarro Calvo A, Martinez-Useros J, Regueira Acebedo N, Fernandez Acenero MJ, Munoz A, et al. Sirt1 mediates the antagonism of Wnt/beta-catenin pathway by vitamin D in colon carcinoma cells. *Int J Biol Sci* 2024;**20**:5495–509.
 86. Li Q, Chan H, Liu WX, Liu CA, Zhou Y, Huang D, et al. Carnobacterium maltaromaticum boosts intestinal vitamin d production to suppress colorectal cancer in female mice. *Cancer Cell* 2023;**41**:1450–65.e8.
 87. Feng YH, Huang YL, Wang Y, Wang P, Song HP, Wang FJ. Antibiotics induced intestinal tight junction barrier dysfunction is associated with microbiota dysbiosis, activated NLRP3 inflammasome and autophagy. *PLoS One* 2019;**14**:e0218384.
 88. Ferreyra JA, Wu KJ, Hryckowian AJ, Bouley DM, Weimer BC, Sonnenburg JL. Gut microbiota-produced succinate promotes c-difficile infection after antibiotic treatment or motility disturbance. *Cell Host Microbe* 2014;**16**:770–7.
 89. Nguyen LH, Ortqvist AK, Cao Y, Simon TG, Roelstraete B, Song MY, et al. Antibiotic use and the development of inflammatory

- bowel disease: a national case-control study in Sweden. *Lancet Gastroenterol* 2020;**5**:986–95.
90. Pull SL, Doherty JM, Mills JC, Gordon JJ, Stappenbeck TS. Activated macrophages are an adaptive element of the colonic epithelial progenitor niche necessary for regenerative responses to injury. *Proc Natl Acad Sci U S A* 2005;**102**:99–104.
 91. Nikolaus S, Schulte B, Al-Massad N, Thieme F, Schulte DM, Bethge J, et al. Increased tryptophan metabolism is associated with activity of inflammatory bowel diseases. *Gastroenterology* 2017;**153**:1504–16.e2.
 92. Liu D, Liang CH, Huang B, Zhuang X, Cui WW, Yang L, et al. Tryptophan metabolism acts as a new anti-ferroptotic pathway to mediate tumor growth. *Adv Sci* 2023;**10**:e2204006.
 93. Michaudel C, Danne C, Agus A, Magniez A, Aucouturier A, Spatz M, et al. Rewiring the altered tryptophan metabolism as a novel therapeutic strategy in inflammatory bowel diseases. *Gut* 2023;**72**:1296–307.
 94. Laurans L, Venteclef N, Haddad Y, Chajadine M, Alzaid F, Metghalchi S, et al. Genetic deficiency of indoleamine 2,3-dioxygenase promotes gut microbiota-mediated metabolic health. *Nat Med* 2018;**24**:1113–20.
 95. Lavoie S, Chun EY, Bae S, Brennan CA, Comeau CAG, Lang JK, et al. Expression of free fatty acid receptor 2 by dendritic cells prevents their expression of interleukin 27 and is required for maintenance of mucosal barrier and immune response against colorectal tumors in mice. *Gastroenterology* 2020;**158**:1359–72.e9.
 96. Wang RX, Lee JS, Campbell EL, Colgan SP. Microbiota-derived butyrate dynamically regulates intestinal homeostasis through regulation of actin-associated protein synaptopodin. *Proc Natl Acad Sci U S A* 2020;**117**:11648–57.
 97. Kim S, Shin YC, Kim TY, Kim Y, Lee YS, Lee SH, et al. Mucin degrader *Akkermansia muciniphila* accelerates intestinal stem cell-mediated epithelial development. *Gut Microbes* 2021;**13**:1–20.
 98. Gutierrez Lopez DE, Lashinger LM, Weinstock GM, Bray MS. Circadian rhythms and the gut microbiome synchronize the host's metabolic response to diet. *Cell Metab* 2021;**33**:873–87.
 99. Tuganbaev T, Mor U, Bashiardes S, Liwinski T, Nobs SP, Leshem A, et al. Diet diurnally regulates small intestinal microbiome-epithelial-immune homeostasis and enteritis. *Cell* 2020;**182**:1441–59.e21.
 100. Talbot J, Hahn P, Kroehling L, Nguyen H, Li D, Littman DR. Feeding-dependent vip neuron-ilc3 circuit regulates the intestinal barrier. *Nature* 2020;**579**:575–80.
 101. Guo X, Wang J, Xu H, Wang Y, Cao Y, Wen Y, et al. Obesity induced disruption on diurnal rhythm of insulin sensitivity via gut microbiome-bile acid metabolism. *Biochim Biophys Acta Mol Cell Biol Lipids* 2024;**1869**:159419.
 102. Choi H, Rao MC, Chang EB. Gut microbiota as a transducer of dietary cues to regulate host circadian rhythms and metabolism. *Nat Rev Gastroenterol Hepatol* 2021;**18**:679–89.
 103. Fawad JA, Luzader DH, Hanson GF, Moutinho Jr TJ, McKinney CA, Mitchell PG, et al. Histone deacetylase inhibition by gut microbe-generated short-chain fatty acids entrains intestinal epithelial circadian rhythms. *Gastroenterology* 2022;**163**:1377–90.e11.
 104. Brooks 2nd JF, Behrendt CL, Ruhn KA, Lee S, Raj P, Takahashi JS, et al. The microbiota coordinates diurnal rhythms in innate immunity with the circadian clock. *Cell* 2021;**184**:4154–67.e12.
 105. Christ A, Lauterbach M, Latz E. Western diet and the immune system: an inflammatory connection. *Immunity* 2019;**51**:794–811.
 106. Zheng M, Zhai Y, Yu Y, Shen J, Chu S, Focaccia E, et al. Tnf compromises intestinal bile-acid tolerance dictating colitis progression and limited infliximab response. *Cell Metab* 2024;**36**:2086–103.e9.
 107. Liu TC, Kern JT, Jain U, Sonnek NM, Xiong S, Simpson KF, et al. Western diet induces paneth cell defects through microbiome alterations and farnesoid X receptor and type i interferon activation. *Cell Host Microbe* 2021;**29**:988–1001.e6.
 108. Yang J, Wei H, Zhou Y, Szeto CH, Li C, Lin Y, et al. High-fat diet promotes colorectal tumorigenesis through modulating gut microbiota and metabolites. *Gastroenterology* 2022;**162**:135–49.e2.
 109. Febbraio MA, Karin M. “Sweet death”: fructose as a metabolic toxin that targets the gut–liver axis. *Cell Metab* 2021;**33**:2316–28.
 110. Thaiss CA, Levy M, Grosheva I, Zheng D, Soffer E, Blacher E, et al. Hyperglycemia drives intestinal barrier dysfunction and risk for enteric infection. *Science* 2018;**359**:1376–83.
 111. Willing BP, Russell SL, Finlay BB. Shifting the balance: antibiotic effects on host-microbiota mutualism. *Nat Rev Microbiol* 2011;**9**:233–43.
 112. Patel VC, Lee S, McPhail MJW, Da Silva K, Guilly S, Zamillo A, et al. Rifaximin-alpha reduces gut-derived inflammation and mucin degradation in cirrhosis and encephalopathy: Rifsys randomised controlled trial. *J Hepatol* 2022;**76**:332–42.
 113. Sawaed J, Zelik L, Levin Y, Feeney R, Naama M, Gordon A, et al. Antibiotics damage the colonic mucus barrier in a microbiota-independent manner. *Sci Adv* 2024;**10**:eadp4119.
 114. Xu D, Gao J, Gilliland 3rd M, Wu X, Song I, Kao JY, et al. Rifaximin alters intestinal bacteria and prevents stress-induced gut inflammation and visceral hyperalgesia in rats. *Gastroenterology* 2014;**146**:484–96.e4.
 115. Li T, Ding N, Guo H, Hua R, Lin Z, Tian H, et al. A gut microbiota-bile acid axis promotes intestinal homeostasis upon aspirin-mediated damage. *Cell Host Microbe* 2024;**32**:191–208.e9.
 116. Rodrigues D, Souza T, Jennen DGJ, Lemmens L, Kleijnans JCS, de Kok TM. Drug-induced gene expression profile changes in relation to intestinal toxicity: state-of-the-art and new approaches. *Cancer Treat Rev* 2019;**77**:57–66.
 117. Soularue E, Lepage P, Colombel JF, Coutzac C, Faleck D, Marthey L, et al. Enterocolitis due to immune checkpoint inhibitors: a systematic review. *Gut* 2018;**67**:2056–67.
 118. Wu D, Hu LN, Han MW, Deng YC, Zhang YM, Ren GQ, et al. PD-1 signaling facilitates activation of lymphoid tissue inducer cells by restraining fatty acid oxidation. *Nat Metab* 2022;**4**:867–82.
 119. Khoshbin K, Khanna L, Maselli D, Atieh J, Breen-Lyles M, Arndt K, et al. Development and validation of test for “leaky gut” small intestinal and colonic permeability using sugars in healthy adults. *Gastroenterology* 2021;**161**:463–75.e13.
 120. Grover M, Camilleri M, Hines J, Burton D, Ryks M, Wadhwa A, et al. C mannitol as a novel biomarker for measurement of intestinal permeability. *Neurogastroenterol Motil* 2016;**28**:1114–9.
 121. Lee GO, McCormick BJJ, Seidman JC, Kosek MN, Haque R, Olortegui MP, et al. Infant nutritional status, feeding practices, enteropathogen exposure, socioeconomic status, and illness are associated with gut barrier function as assessed by the lactulose mannitol test in the mal-ed birth cohort. *Am J Trop Med Hyg* 2017;**97**:281–90.
 122. Olivera PA, Martinez Lozano H, Leibovitz H, Xue M, Neustaeter A, Espin Garcia O, et al. Healthy first-degree relatives from multiplex families vs simplex families have higher subclinical intestinal inflammation, a distinct fecal microbial signature, and harbor a higher risk of developing Crohn's disease. *Gastroenterology* 2025;**168**:99–110.e2.
 123. Turpin W, Lee SH, Garay JAR, Madsen KL, Meddings JB, Bedrani L, et al. Increased intestinal permeability is associated with later development of Crohn's disease. *Gastroenterology* 2020;**159**:2092–100.e5.
 124. Sorribas M, Jakob MO, Yilmaz B, Li H, Stutz D, Noser Y, et al. FXR modulates the gut–vascular barrier by regulating the entry sites for bacterial translocation in experimental cirrhosis. *J Hepatol* 2019;**71**:1126–40.
 125. Demaude J, Salvador-Cartier C, Fioramonti J, Ferrier L, Bueno L. Phenotypic changes in colonocytes following acute stress or activation of mast cells in mice: implications for delayed epithelial barrier dysfunction. *Gut* 2006;**55**:655–61.

126. D'Agostino L, Daniele B, Pignata S, Barone MV, Ciacci C, Sollazzo R, et al. Postheparin plasma diamine oxidase increases in patients with coeliac disease during gluten free diet. *Gut* 1987;**28**(Suppl):131–4.
127. D'Agostino L, Daniele B, Pallone F, Pignata S, Leoni M, Mazzacca G. Postheparin plasma diamine oxidase in patients with small bowel Crohn's disease. *Gastroenterology* 1988;**95**:1503–9.
128. Cao B, Zhao RY, Li HH, Xu XM, Cui H, Deng H, et al. Oral administration of asparagine and 3-indolepropionic acid prolongs survival time of rats with traumatic colon injury. *Mil Med Res* 2022;**9**:37.
129. Playford RJ, Belo A, Poulson R, Fitzgerald AJ, Harris K, Pawluczyk I, et al. Effects of mouse and human lipocalin homologues 24p3/lcn2 and neutrophil gelatinase-associated lipocalin on gastrointestinal mucosal integrity and repair. *Gastroenterology* 2006;**131**:809–17.
130. Xia MY, Lu JM, Lan JB, Teng T, Shiao R, Sun HB, et al. Elevated IL-22 as a result of stress-induced gut leakage suppresses septal neuron activation to ameliorate anxiety-like behavior. *Immunity* 2025;**58**:218–31.e12.
131. Tulkens J, Vergauwen G, Van Deun J, Geeurickx E, Dhondt B, Lippens L, et al. Increased levels of systemic lps-positive bacterial extracellular vesicles in patients with intestinal barrier dysfunction. *Gut* 2020;**69**:191–3.
132. Schmitz H, Barmeyer C, Fromm M, Runkel N, Foss HD, Bentzel CJ, et al. Altered tight junction structure contributes to the impaired epithelial barrier function in ulcerative colitis. *Gastroenterology* 1999;**116**:301–9.
133. Graham WV, Wang F, Clayburgh DR, Cheng JX, Yoon B, Wang Y, et al. Tumor necrosis factor-induced long myosin light chain kinase transcription is regulated by differentiation-dependent signaling events. Characterization of the human long myosin light chain kinase promoter. *J Biol Chem* 2006;**281**:26205–15.
134. Heller F, Florian P, Bojarski C, Richter J, Christ M, Hillenbrand B, et al. Interleukin-13 is the key effector TH2 cytokine in ulcerative colitis that affects epithelial tight junctions, apoptosis, and cell restitution. *Gastroenterology* 2005;**129**:550–64.
135. Oami T, Abtahi S, Shimazui T, Chen CW, Sweat YY, Liang Z, et al. Claudin-2 upregulation enhances intestinal permeability, immune activation, dysbiosis, and mortality in sepsis. *Proc Natl Acad Sci U S A* 2024;**121**:e2217877121.
136. Ma Y, Yue J, Zhang Y, Shi C, Odenwald M, Liang WG, et al. Acf7 regulates inflammatory colitis and intestinal wound response by orchestrating tight junction dynamics. *Nat Commun* 2017;**8**:15375.
137. Hollander D, Vadheim CM, Brettholz E, Petersen GM, Delahunty T, Rotter JI. Increased intestinal permeability in patients with Crohn's disease and their relatives. A possible etiologic factor. *Ann Intern Med* 1986;**105**:883–5.
138. Waehrens R, Ohlsson H, Sundquist J, Sundquist K, Zoller B. Risk of irritable bowel syndrome in first-degree, second-degree and third-degree relatives of affected individuals: a nationwide family study in Sweden. *Gut* 2015;**64**:215–21.
139. Schmitt M, Greten FR. The inflammatory pathogenesis of colorectal cancer. *Nat Rev Immunol* 2021;**21**:653–67.
140. Grivennikov SI, Wang K, Mucida D, Stewart CA, Schnabl B, Jauch D, et al. Adenoma-linked barrier defects and microbial products drive IL-23/IL-17-mediated tumour growth. *Nature* 2012;**491**:254–8.
141. Pu Y, Song Y, Zhang MD, Long CF, Li J, Wang YA, et al. Golm1 restricts colitis and colon tumorigenesis by ensuring notch signaling equilibrium in intestinal homeostasis. *Signal Transduct Tar* 2021;**6**:148.
142. Wei M, Ma Y, Shen L, Xu Y, Liu L, Bu X, et al. NDRG2 regulates adherens junction integrity to restrict colitis and tumorigenesis. *EBioMedicine* 2020;**61**:103068.
143. Tilg H, Adolph TE, Trauner M. Gut–liver axis: pathophysiological concepts and clinical implications. *Cell Metab* 2022;**34**:1700–18.
144. Gabele E, Dostert K, Hofmann C, Wiest R, Scholmerich J, Hellerbrand C, et al. DSS induced colitis increases portal LPS levels and enhances hepatic inflammation and fibrogenesis in experimental NASH. *J Hepatol* 2011;**55**:1391–9.
145. Zheng C, Wang L, Zou T, Lian S, Luo J, Lu Y, et al. Ileitis promotes masld progression via bile acid modulation and enhanced TGR5 signaling in ileal CD8⁺ T cells. *J Hepatol* 2024;**80**:764–77.
146. Loomba R, Seguritan V, Li W, Long T, Klitgord N, Bhatt A, et al. Gut microbiome-based metagenomic signature for non-invasive detection of advanced fibrosis in human nonalcoholic fatty liver disease. *Cell Metab* 2017;**25**:1054–62.e5.
147. Fizanne L, Villard A, Benabbou N, Recoquillon S, Soletti R, Delage E, et al. Faeces-derived extracellular vesicles participate in the onset of barrier dysfunction leading to liver diseases. *J Extracell Vesicles* 2023;**12**:e12303.
148. Rungratanawanich W, Lin YH, Wang X, Kawamoto T, Chidambaram SB, Song BJ. ALDH2 deficiency increases susceptibility to binge alcohol-induced gut leakiness, endotoxemia, and acute liver injury in mice through the gut–liver axis. *Redox Biol* 2023;**59**:102577.
149. Hao LY, Zhong W, Wei XY, Woo JM, Dong HB, Guo W, et al. Intestinal conventional type 1 dendritic cell protects against alcohol-induced steatohepatitis by maintaining *Akkermansia muciniphila* abundance. *Hepatology* 2022;**76**:S984.
150. Kummen M, Holm K, Anmarkrud JA, Nygard S, Vesterhus M, Hoivik ML, et al. The gut microbial profile in patients with primary sclerosing cholangitis is distinct from patients with ulcerative colitis without biliary disease and healthy controls. *Gut* 2017;**66**:611–9.
151. Nakamoto N, Sasaki N, Aoki R, Miyamoto K, Suda W, Teratani T, et al. Gut pathobionts underlie intestinal barrier dysfunction and liver t helper 17 cell immune response in primary sclerosing cholangitis. *Nat Microbiol* 2019;**4**:492–503.
152. Liu R, Li X, Huang Z, Zhao D, Ganesh BS, Lai G, et al. C/EBP homologous protein-induced loss of intestinal epithelial stemness contributes to bile duct ligation-induced cholestatic liver injury in mice. *Hepatology* 2018;**67**:1441–57.
153. Gui W, Hole MJ, Molinaro A, Edlund K, Jorgensen KK, Su H, et al. Colitis ameliorates cholestatic liver disease via suppression of bile acid synthesis. *Nat Commun* 2023;**14**:3304.
154. Liao L, Schneider KM, Galvez EJC, Frissen M, Marschall HU, Su H, et al. Intestinal dysbiosis augments liver disease progression via NLRP3 in a murine model of primary sclerosing cholangitis. *Gut* 2019;**68**:1477–92.
155. Su L, Nalle SC, Shen L, Turner ES, Singh G, Breskin LA, et al. Tnfr2 activates MLCK-dependent tight junction dysregulation to cause apoptosis-mediated barrier loss and experimental colitis. *Gastroenterology* 2013;**145**:407–15.
156. Odenwald MA, Turner JR. Intestinal permeability defects: is it time to treat?. *Clin Gastroenterol Hepatol* 2013;**11**:1075–83.
157. Marchelletta RR, Krishnan M, Spalinger MR, Placone TW, Alvarez R, Sayoc-Becerra A, et al. T cell protein tyrosine phosphatase protects intestinal barrier function by restricting epithelial tight junction remodeling. *J Clin Invest* 2021;**131**:e138230.
158. Wu H, Xie S, Miao J, Li Y, Wang Z, Wang M, et al. Lactobacillus reuteri maintains intestinal epithelial regeneration and repairs damaged intestinal mucosa. *Gut Microbes* 2020;**11**:997–1014.
159. Seo B, Jeon K, Moon S, Lee K, Kim WK, Jeong H, et al. *Roseburia* spp. Abundance associates with alcohol consumption in humans and its administration ameliorates alcoholic fatty liver in mice. *Cell Host Microbe* 2020;**27**:25–40.e6.
160. Zhang Y, Wang Y, Zhang X, Wang P, Shi F, Zhang Z, et al. Gastrointestinal self-adaptive and nutrient self-sufficient *Akkermansia muciniphila*-gelatin porous microgels for synergistic therapy of ulcerative colitis. *ACS Nano* 2024;**18**:26807–27.
161. Chen Y, Lin S, Wang L, Zhang Y, Chen H, Fu Z, et al. Reinforcement of the intestinal mucosal barrier via mucus-penetrating pegylated bacteria. *Nat Biomed Eng* 2024:823–41.
162. Cao F, Jin L, Gao Y, Ding Y, Wen H, Qian Z, et al. Artificial-enzymes-armed bifidobacterium longum probiotics for alleviating

- intestinal inflammation and microbiota dysbiosis. *Nat Nanotechnol* 2023;**18**:617–27.
163. Perino A, Demagny H, Velazquez-Villegas L, Schoonjans K. Molecular physiology of bile acid signaling in health, disease, and aging. *Physiol Rev* 2021;**101**:683–731.
 164. Friedman ES, Li Y, Shen TCD, Jiang J, Chau L, Adorini L, et al. FXR-dependent modulation of the human small intestinal microbiome by the bile acid derivative obeticholic acid. *Gastroenterology* 2018;**155**:1741–52.e5.
 165. Mouries J, Brescia P, Silvestri A, Spadoni I, Sorribas M, Wiest R, et al. Microbiota-driven gut vascular barrier disruption is a prerequisite for non-alcoholic steatohepatitis development. *J Hepatol* 2019;**71**:1216–28.
 166. Zhou XL, Chakraborty D, Murray IA, Coslo D, Kehs Z, Vijay A, et al. Aryl hydrocarbon receptor activation coordinates mouse small intestinal epithelial cell programming. *Lab Invest* 2023;**103**:100012.
 167. Hu XC, Xiao WF, Lei YX, Green A, Lee X, Maradana MR, et al. Aryl hydrocarbon receptor utilises cellular zinc signals to maintain the gut epithelial barrier. *Nat Commun* 2023;**14**:5431.
 168. Singh R, Chandrashekarappa S, Bodduluri SR, Baby BV, Hegde B, Kotla NG, et al. Enhancement of the gut barrier integrity by a microbial metabolite through the Nrf2 pathway. *Nat Commun* 2019;**10**:89.
 169. Wrzosek L, Ciocan D, Hugot C, Spatz M, Dupeux M, Houron C, et al. Microbiota tryptophan metabolism induces aryl hydrocarbon receptor activation and improves alcohol-induced liver injury. *Gut* 2021;**70**:1299–308.
 170. Qian M, Liu J, Zhao D, Cai P, Pan C, Jia W, et al. Aryl hydrocarbon receptor deficiency in intestinal epithelial cells aggravates alcohol-related liver disease. *Cell Mol Gastroenterol Hepatol* 2022;**13**:233–56.
 171. Xue C, Li G, Zheng Q, Gu X, Shi Q, Su Y, et al. Tryptophan metabolism in health and disease. *Cell Metab* 2023;**35**:1304–26.
 172. Wu P, Yao S, Wang X, Yang L, Wang S, Dai W, et al. Oral administration of nanoformulated indoximod ameliorates ulcerative colitis by promoting mitochondrial function and mucosal healing. *Int J Pharm* 2023;**637**:122813.
 173. Wang D, Li D, Zhang Y, Chen J, Zhang Y, Liao C, et al. Functional metabolomics reveal the role of AHR/GPR35 mediated kynurenic acid gradient sensing in chemotherapy-induced intestinal damage. *Acta Pharm Sin B* 2021;**11**:763–80.
 174. Wang G, Fan Y, Zhang G, Cai S, Ma Y, Yang L, et al. Microbiota-derived indoles alleviate intestinal inflammation and modulate microbiome by microbial cross-feeding. *Microbiome* 2024;**12**:59.
 175. Zhang Y, Liu R, Chen Y, Cao Z, Liu C, Bao R, et al. *Akkermansia muciniphila* supplementation in patients with overweight/obese type 2 diabetes: efficacy depends on its baseline levels in the gut. *Cell Metab* 2025;**37**:592–605.e6.
 176. Lamas B, Richard ML, Leducq V, Pham HP, Michel ML, Da Costa G, et al. CARD9 impacts colitis by altering gut microbiota metabolism of tryptophan into aryl hydrocarbon receptor ligands. *Nat Med* 2016;**22**:598–605.
 177. Guo P, Wang W, Xiang Q, Pan C, Qiu Y, Li T, et al. Engineered probiotic ameliorates ulcerative colitis by restoring gut microbiota and redox homeostasis. *Cell Host Microbe* 2024;**32**:1502–18.e9.
 178. Santos AJM, Lo YH, Mah AT, Kuo CJ. The intestinal stem cell niche: homeostasis and adaptations. *Trends Cell Biol* 2018;**28**:1062–78.
 179. Jee J, Park JH, Im JH, Kim MS, Park E, Lim T, et al. Functional recovery by colon organoid transplantation in a mouse model of radiation proctitis. *Biomaterials* 2021;**275**:120925.
 180. Sugimoto S, Kobayashi E, Fujii M, Ohta Y, Arai K, Matano M, et al. An organoid-based organ-repurposing approach to treat short bowel syndrome. *Nature* 2021;**592**:99–104.
 181. Danese S, Vermeire S, Hellstern P, Panaccione R, Rogler G, Fraser G, et al. Randomised trial and open-label extension study of an anti-interleukin-6 antibody in Crohn's disease (andante I and II). *Gut* 2019;**68**:40–8.
 182. Ouyang W, O'Garra A. IL-10 family cytokines IL-10 and IL-22: from basic science to clinical translation. *Immunity* 2019;**50**:871–91.
 183. Liu J, Ren H, Zhang C, Li J, Qiu Q, Zhang N, et al. Orally-delivered, cytokine-engineered extracellular vesicles for targeted treatment of inflammatory bowel disease. *Small* 2023;**19**:e2304023.
 184. Zegarra Ruiz DF, Kim DV, Norwood K, Saldana-Morales FB, Kim M, Ng C, et al. Microbiota manipulation to increase macrophage IL-10 improves colitis and limits colitis-associated colorectal cancer. *Gut Microbes* 2022;**14**:2119054.
 185. He GW, Lin L, DeMartino J, Zheng X, Staliarova N, Dayton T, et al. Optimized human intestinal organoid model reveals interleukin-22-dependency of paneth cell formation. *Cell Stem Cell* 2022;**29**:1718–20.
 186. Chiang HY, Lu HH, Sudhakar JN, Chen YW, Shih NS, Weng YT, et al. IL-22 initiates an IL-18-dependent epithelial response circuit to enforce intestinal host defence. *Nat Commun* 2022;**13**:874.
 187. Wagner F, Mansfield JC, Lekkerkerker AN, Wang Y, Keir M, Dash A, et al. Dose escalation randomised study of efmarodocokin alfa in healthy volunteers and patients with ulcerative colitis. *Gut* 2023;**72**:1451–61.
 188. Gaudino SJ, Singh A, Huang H, Padiadpu J, Jean Pierre M, Kempen C, et al. Intestinal IL-22RA1 signaling regulates intrinsic and systemic lipid and glucose metabolism to alleviate obesity-associated disorders. *Nat Commun* 2024;**15**:1597.
 189. Zhang P, Liu J, Lee A, Tsaur I, Ohira M, Duong V, et al. IL-22 resolves masld via enterocyte Stat3 restoration of diet-perturbed intestinal homeostasis. *Cell Metab* 2024;**36**:2341–54.e6.
 190. Lin X, Gaudino SJ, Jang KK, Bahadur T, Singh A, Banerjee A, et al. IL-17RA-signaling in Lgr5⁺ intestinal stem cells induces expression of transcription factor atoh1 to promote secretory cell lineage commitment. *Immunity* 2022;**55**:237–53.e8.
 191. Na YR, Jung D, Stakenborg M, Jang H, Gu GJ, Jeong MR, et al. Prostaglandin E2 receptor PTGER4-expressing macrophages promote intestinal epithelial barrier regeneration upon inflammation. *Gut* 2021;**70**:2249–60.
 192. Fu Y, Ding X, Zhang M, Feng C, Yan Z, Wang F, et al. Intestinal mucosal barrier repair and immune regulation with an AI-developed gut-restricted PHD inhibitor. *Nat Biotechnol* 2024. <https://doi.org/10.1038/s41587-024-02503-w>. Available from: <https://doi.org/10.1038/s41587-024-02503-w>.
 193. Yang J, Yang X, Wu G, Huang F, Shi X, Wei W, et al. Gut microbiota modulate distal symmetric polyneuropathy in patients with diabetes. *Cell Metab* 2023;**35**:1548–62.e7.
 194. Liu J, MacNaughtan J, Kerbert AJC, Portlock T, Martinez Gonzalez J, Jin Y, et al. Clinical, experimental and pathophysiological effects of YAQ-001: a non-absorbable, gut-restricted adsorbent in models and patients with cirrhosis. *Gut* 2024;**73**:1183–98.
 195. Vazquez Roque MI, Camilleri M, Smyrk T, Murray JA, Marietta E, O'Neill J, et al. A controlled trial of gluten-free diet in patients with irritable bowel syndrome-diarrhea: effects on bowel frequency and intestinal function. *Gastroenterology* 2013;**144**:903–11.e3.
 196. Li H, Zhang L, Li J, Wu Q, Qian L, He J, et al. Resistant starch intake facilitates weight loss in humans by reshaping the gut microbiota. *Nat Metab* 2024;**6**:578–97.
 197. Levine A, Wine E, Assa A, Sigall Boneh R, Shaoul R, Kori M, et al. Crohn's disease exclusion diet plus partial enteral nutrition induces sustained remission in a randomized controlled trial. *Gastroenterology* 2019;**157**:440–50.e8.
 198. Zhou Q, Verne ML, Fields JZ, Lefante JJ, Basra S, Salameh H, et al. Randomised placebo-controlled trial of dietary glutamine supplements for postinfectious irritable bowel syndrome. *Gut* 2019;**68**:996–1002.
 199. Schuppan D, Maki M, Lundin KEA, Isola J, Friesing-Sosnik T, Taavela J, et al. A randomized trial of a transglutaminase 2 inhibitor for celiac disease. *N Engl J Med* 2021;**385**:35–45.
 200. Dotsenko V, Tewes B, Hils M, Pasternack R, Isola J, Taavela J, et al. Transcriptomic analysis of intestine following administration of a

- transglutaminase 2 inhibitor to prevent gluten-induced intestinal damage in celiac disease. *Nat Immunol* 2024;**25**:1218–30.
201. Kotla NG, Rochev Y. IBD disease-modifying therapies: insights from emerging therapeutics. *Trends Mol Med* 2023;**29**:241–53.
 202. Luissint AC, Parkos CA, Nusrat A. Inflammation and the intestinal barrier: leukocyte-epithelial cell interactions, cell junction remodeling, and mucosal repair. *Gastroenterology* 2016;**151**:616–32.
 203. Rinella ME, Lazarus JV, Ratziu V, Francque SM, Sanyal AJ, Kanwal F, et al. A multisociety delphi consensus statement on new fatty liver disease nomenclature. *J Hepatol* 2023;**79**:1542–56.
 204. Lee BP, Dodge JL, Terrault NA. National prevalence estimates for steatotic liver disease and subclassifications using consensus nomenclature. *Hepatology* 2024;**79**:666–73.
 205. Lin A, Roth H, Anyane-Yeboah A, Rubin DT, Paul S. Prevalence of nonalcoholic fatty liver disease in patients with inflammatory bowel disease: a systematic review and meta-analysis. *Inflamm Bowel Dis* 2021;**27**:947–55.
 206. Kuang J, Wang J, Li Y, Li M, Zhao M, Ge K, et al. Hydoxycholeic acid alleviates non-alcoholic fatty liver disease through modulating the gut–liver axis. *Cell Metab* 2023;**35**:1752–66.e8.
 207. Jiang J, Ma Y, Liu Y, Lu D, Gao X, Krausz KW, et al. Glycine-beta-muricholic acid antagonizes the intestinal farnesoid X receptor–ceramide axis and ameliorates NASH in mice. *Hepatol Commun* 2022;**6**:3363–78.
 208. Silvestri A, Gil Gomez A, Vitale M, Braga D, Demitri C, Brescia P, et al. Biomimetic superabsorbent hydrogel acts as a gut protective dynamic exoskeleton improving metabolic parameters and expanding a. *Muciniphila*. *Cell Rep Med* 2023;**4**:101235.
 209. Hao L, Zhong W, Woo J, Wei X, Ma H, Dong H, et al. Conventional type 1 dendritic cells protect against gut barrier disruption via maintaining *Akkermansia muciniphila* in alcoholic steatohepatitis. *Hepatology* 2023;**78**:896–910.